



# **Source Water Assessment**

A Hydrogeologic Susceptibility and
Vulnerability Assessment for
ConocoPhillips
Kuparuk Operations Center (KOC)

PWSID # 330031.001

July 2004

Drinking Water Protection Program Report #1537 Alaska Department of Environmental Conservation

# Source Water Assessment for the ConocoPhillips Kuparuk Operations Center (KOC)

PWSID# 330031.001

July 2004

Drinking Water Protection Program Report #1537

The Drinking Water Protection Program (DWPP) is producing Source Water Assessments in compliance with the Safe Drinking Water Act Amendments of 1996. Each assessment includes a delineation of the source water area, an inventory of potential and existing contaminant sources that may impact the water, a risk ranking for each of these contaminants, and an evaluation of the potential vulnerability of these drinking water sources.

These assessments are intended to provide public water systems owners/operators, communities, and local governments with the best available information that may be used to protect the quality of their drinking water. The assessments combine information obtained from various sources, including the U.S. Environmental Protection Agency, Alaska Department of Environmental Conservation (ADEC), public water system owners/operators, and other public information sources. The results of this assessment are subject to change if additional data becomes available. It is anticipated this assessment will be updated every five years to reflect any changes in the vulnerability and/or susceptibility of public drinking water source. If you have any additional information that may affect the results of this assessment, please contact the Program Coordinator of DWPP, (907) 269-7521.

# **CONTENTS**

			Page
SECTION	Drin Kup Inve Ranl Vulr	cutive Summary Iking Water System and Area Overview aruk Operations Center Drinking Water Protection Area entory of Potential and Existing Contaminant Sources king of Contaminant Risks herability of the Drinking Water System erences	1 1 2 2 2 7
		TABLES	
TABLE	2. 3.	Definition of Zones Susceptibility of the Water Source Contaminant Risks Overall Vulnerability	1 3 3 3
		APPENDICES	
APPENDIX	A.	Kuparuk Operations Center Drinking Water Protection Area (Map 1)	
	B.	Contaminant Source Inventory and Risk Rankings (Tables 1 - 4)	
	C.	Kuparuk Operations Center Drinking Water Protection Area and Potential and Existing Contaminant Sources (Map 2)	
	D.	Vulnerability Analysis and Contaminant Risks (Charts $1-13$ )	

# **Drinking Water Protection Program Alaska Department of Environmental Conservation**

#### **EXECUTIVE SUMMARY**

The ConocoPhillips - Kuparuk Operation Center (KOC) water system is a Class A (community) water system that obtains water from a small reservoir that receives water from the Ugnuravik River. The intake is located approximately 0.5-miles west of the Kuparuk base camp. Access to the intake area is not restricted. The overall protection area for the intake is approximately 76 acres in size and received a susceptibility rating of "high". A rating of high to very high is typical for all systems with surface water intakes. Potential and existing sources of the following contaminants were evaluated for the Source Water Assessment: bacteria and viruses, nitrates and/or nitrites, heavy metals, cyanide, and other inorganic chemicals, synthetic organic chemicals, volatile organic chemicals, and other organic chemicals. An oil pipeline was identified as a potential source of contaminants for the drinking water source. This evaluation included all available water sampling data submitted to ADEC by the system operator. The samples may have been collected from either raw water or post-treated water. Combining the susceptibility of the surface water source with the contaminant risks, this water system has received a vulnerability rating of "low" for bacteria and nitrates/nitrites, and synthetic organic chemicals; "medium" for heavy metals; and "high" for volatile organic chemicals and other organic chemicals. This assessment can be used as a foundation for local voluntary protection efforts as well as a basis for the continuous efforts on the part of ConocoPhillips to protect public health.

# DRINKING WATER SYSTEM AND AREA OVERVIEW

The Kuparuk Operations Center (KOC) (Sec. 9, T011N, R010E, Umiat Meridian) is located approximately 27 miles west of Prudhoe Bay along the Ugnuravik River (Please see the inset of Map 1 in Appendix A for location). The water system is a Class A (community) water system that serves the working population year-round and obtains water from a small reservoir that receives water from the Ugnuravik River. The intake is located approximately 0.5-miles west of the Kuparuk base camp (See Map 1 of Appendix A).

Summer temperatures in the Kuparuk area range from the 20s to the 70s; winter temperatures average -40 to

10. Average annual precipitation is approximately 5 inches, including 30 inches of snowfall (ADCED, 2004).

The 1998 sanitary survey indicates that intake is screened, maintained, and protected from ice buildup and siltation. The survey also states that the pump capacity of the system is 300 gpm with 40,000 gallons of storage capacity. System operators report that the reservoir has a capacity of approximately 299 acre-feet and that the Ugnaravik River generally flows at less than 10 cfs during the summer months.

# KUPARUK OPERATIONS CENTER DRINKING WATER PROTECTION AREA

Identifying the pathways most likely for surface contamination to reach water intake areas is the first step in determining the water system's risk. These are initially determined by looking at the drainage area contributing overland water flow to a surface water source intake. The entire drainage area is also known as the "drinking water protection area". Please refer to pages 10-11of the "Guidance Manual for Class A Public Water Systems" for additional information.

The protection area established for surface water sources by the ADEC is usually separated into three zones, limited by the watershed boundary. These zones correspond to the overland-flow distance that water travels to get to the source. The ADEC Drinking Water Protection Program's Technical Advisory Committee developed guidelines for derivation of these zones in 1998. The following is a summary of the three protection area zones:

Table 1. Definition of Zones

Zone	Definition
A	Areas within 1000-ft of lakes or streams
В	Areas within 1-mile of lakes or streams
C	The watershed boundary

The protection area for KOC intake includes each of these Zones (See Map 1 of Appendix A).

# INVENTORY OF POTENTIAL AND EXISTING CONTAMINANT SOURCES

The Drinking Water Protection Program has completed an inventory of potential and existing sources of contamination within the KOC protection area. This inventory was completed through a search of agency records and other publicly available information. There is a wide array of potential contamination sources to surface water. These contaminants are found within agricultural, residential, commercial, and industrial areas, but *can also occur within areas that have little or no development*.

For Class A public water system assessments, six categories of drinking water contaminants were inventoried. They include:

- Bacteria and viruses:
- Nitrates and/or nitrites;
- Volatile organic chemicals;
- Heavy metals, cyanide, and other inorganic chemicals;
- Synthetic Organic Chemicals; and
- Other Organic Chemicals.

Sources identified in the protection area are displayed on Map 2 of Appendix C and summarized in Table 1 of Appendix B.

#### RANKING OF CONTAMINANT RISKS

Once potential and existing sources of contamination have been identified, they are assigned a ranking according to what category and level of risk they represent. Ranking of contaminant risks for "potential" or "existing" sources of contamination is a function of the toxicity and the volume of specific contaminants associated with that source. Rankings include:

- Low;
- Medium;
- High; and
- Very High.

The time-of-travel for contaminants within the water is dependent on the physical and chemical characteristics of each contaminant. Bacteria and Viruses are only inventoried in Zone A because of their short life span. Only "Very High" and "High" rankings are inventoried within Zones B and C due to the probability of contaminant dilution by the time the contaminants reach the water intake.

The remaining tables in Appendix B (if necessary) contain the ranking of potential and existing sources of

contamination with respect to bacteria and viruses, nitrates and/or nitrites, volatile organic chemicals, heavy metals, cyanide, and other inorganic chemicals, synthetic organic chemicals, and other organic chemicals.

# VULNERABILITY OF THE DRINKING WATER SYSTEM

Vulnerability of a drinking water source to contamination is a combination of two factors:

- Surface Water Susceptibility; and
- Contaminant risks.

Appendix D contains 13 charts, which together form the 'Vulnerability Analysis' for the public drinking water Source Water Assessment. Chart 1 analyzes the 'Susceptibility of the Surface Water Source' to contamination by looking at the climate, terrain, and intake location. Chart 2 analyzes 'Contaminant Risks' for the drinking water source with respect to bacteria and viruses. The 'Contaminant Risks' portion of the analysis considers potential sources of contaminants as well as a review of contamination that has or may have occurred, but has not arrived or been detected at the intake area. Chart 3 contains the 'Vulnerability Analysis for Bacteria and Viruses', which is a composite score of the Vulnerability Analysis and the overall Susceptibility. Charts 4 through 13 repeat the Contaminant Risks and Vulnerability Analyses for nitrates and nitrites, volatile organic chemicals, heavy metals, evanide, and other inorganic chemicals. synthetic organic chemicals, and other organic chemicals, respectively.

A score for the Surface Water Susceptibility of the source is reached by considering the properties of the water intake and the surrounding area. The derivation of this information is presented below and the data for this source is shown in Chart 1 of Appendix D.

Susceptibility of the Surface Water Source – always considered to be "high" (30 points)

+

Adequate Construction of the Intake (0-5 Points)

+

Runoff Potential Within Zone B (0 - 5 Points)

+

Dilution Capacity of the Surface Water (0 - 10 Points)

=

Natural Susceptibility (0 – 50 Points)

A ranking is assigned for the Surface Water Susceptibility according to the point score:

Surface Water Source Susceptibility Ratings

40 to 50 pts Very High 30 to < 40 pts High

Table 2. Susceptibility of the Water Source

	Score	Rating
Minimum Allowable	30	
Susceptibility		
Intake Construction	0	
Adequate		
Runoff Potential	0	
Dilution Capacity	5	
Overall Susceptibility	35	High

For contaminants, risks to a drinking water source depend on the type, number or density, and distribution of the contaminant sources. The Contaminant Risk score has been derived from an examination of existing, and historical contamination sources that have been detected in the protection area through routine sampling. It also evaluates potential sources of contamination. Flow charts are used to assign a point score, and ratings are assigned in the same way as the susceptibility:

Contaminant Ri	sk Ratings
40 to 50 pts	Very High
30 to < 40 pts	High
20 to < 30 pts	Medium
< 20 pts	Low

Table 3 summarizes the Contaminant Risks for each category of drinking water contaminants.

**Table 3. Contaminant Risks** 

Category	Score	Rating
Bacteria and Viruses	0	Low
Nitrates and/or Nitrites	0	Low
Volatile Organic Chemicals	25	Medium
Heavy Metals, Cyanide, and		
Other Inorganic Chemicals	12	Low
Synthetic Organic Chemicals	0	Low
Other Organic Chemicals	40	Very High

Finally, an overall vulnerability score is assigned for each contaminant type by combining each of the contaminant risk scores with the susceptibility score:

Susceptibility of the Surface Water Source

(0-50 points)

+

Contaminant Risks (0 - 50 points)

=

Vulnerability of the Drinking Water Source to Contamination (0 - 100).

Again, rankings are assigned according to a point score:

Overall Vulneral	oility Ratings
80 to 100 pts	Very High
60 to < 80 pts	High
40 to < 60 pts	Medium
< 40 pts	Low

Table 4 contains the overall vulnerability scores and ratings for each of the six categories of drinking water contaminants. Note: scores are rounded off to the nearest five.

Table 4. Overall Vulnerability

Category	Score	Rating
Bacteria and Viruses	35	Low
Nitrates and Nitrites	35	Low
Volatile Organic Chemicals	60	High
Heavy Metals, Cyanide, and		
Other Inorganic Chemicals	45	Medium
Synthetic Organic Chemicals	35	Low
Other Organic Chemicals	75	High

#### **Bacteria and Viruses**

The contaminant risk for bacteria and viruses is "low". Typically, coliform detection in raw water samples collected from surface water sources is normal. (See Chart 2 – Contaminant Risks for Bacteria and Viruses in Appendix D).

Coliforms (a bacteria) are found naturally in the environment and although they aren't necessarily a health threat, they are an indicator of other potentially harmful bacteria in the water, more specifically, fecal coliforms and E. coli which only come from human and animal fecal waste. Harmful bacteria can cause diarrhea, cramps, nausea, headaches, or other symptoms (EPA, 2003). Positive samples increase the overall vulnerability of the drinking water source, indicating that the source is susceptible to bacteria and virus contamination.

Positive bacteria counts have not been detected in recent sampling.

After combining the contaminant risk for bacteria and viruses with the natural susceptibility of the source, the overall vulnerability of the source to bacteria and virus contamination is "low".

#### **Nitrates and Nitrites**

The contaminant risk for nitrates and nitrites is "low" (See Chart 4 - Contaminant Risks for Nitrates and/or Nitrites in Appendix D). Nitrates are very mobile, moving at approximately the same rate as water.

Sampling history for the water source indicates that nitrates have not been detected in samples collected in 2000 and 2003. The Maximum Contaminant Level (MCL) for nitrates is 10 milligrams per liter (mg/L). The MCL is the maximum level of contaminant that is allowed to exist in drinking water and still be consumed by humans without harmful health effects (EPA, 2003).

Possible sources of nitrates/nitrites could be from airfields, human/animal activity along roads, ATV trails or snowmachine trails located within the protection area.

After combining the contaminant risk for nitrates and nitrites with the natural susceptibility of the source, the overall vulnerability of the source to contamination is "low".

#### **Volatile Organic Chemicals**

The contaminant risk for volatile organic chemicals is "medium" (See Chart 6 – Contaminant Risks for Volatile Organic Chemicals in Appendix D).

Trihalomethanes and chloroform were detected at levels below the MCL during sampling in 2003-4, although these chemicals typically originate during the process of water treatment and not from the source waters. The MCL for chloroform is 0.2 milligrams per liter (mg/L) and the MCL for total trihalomethanes is 0.1 mg/L.

Possible sources of volatile organic chemicals could be from oil spillage, airfields, human/animal activity along roads, ATV trails or snowmachine trails located within the protection area.

After combining the contaminant risk for volatile organic chemicals with the natural susceptibility of the source, the overall vulnerability of the source to contamination is "high".

# Heavy Metals, Cyanide, and Other Inorganic Chemicals

The contaminant risk for heavy metals is "low". Copper and lead were detected in samples collected during 2000-2002 at levels below the MCL (See Chart 8 – Contaminant Risks for Heavy Metals, Cyanide, and Other Inorganic Chemicals in Appendix D). The MCL for copper is 1.3 mg/l. and the MCL for lead is 0.015 mg/l.

The most common source of these chemicals is the infrastructure of the distribution system following the treatment process and not from the source waters. Additional sources of heavy metals could be from oil spillage, airfields, activity along roads, ATV or snowmachine trails located within the protection area.

After combining the contaminant risk for heavy metals with the natural susceptibility of the source, the overall vulnerability of the well to contamination is "medium".

#### **Synthetic Organic Chemicals**

The contaminant risk for synthetic organic chemicals is "low".

Review of the historical sampling data indicates that test results for ethylene dibromide in 2003 were negative.

After combining the contaminant risk with the natural susceptibility of the source, the overall vulnerability to synthetic organic chemicals of the source is "low" (See Chart 11 – Contaminant Risks for Synthetic Organic Chemicals in Appendix D).

#### **Other Organic Chemicals**

The contaminant risk for other organic chemicals is "very high".

Review of the historical sampling data indicates that no other organic chemicals have been sampled recently.

A possible source of other organic chemicals could be from oil spillage within the protection area.

After combining the contaminant risk with the natural susceptibility of the source, the overall vulnerability to other organic chemicals of the source is "high" (See Chart 13 – Contaminant Risks for Other Organic Chemicals in Appendix D).

#### **Using the Source Water Assessment**

This assessment of contaminant risks can be used as a foundation for local voluntary protection efforts as well as a basis for the continuous efforts on the part of ConocoPhillips to protect public health. It is anticipated that Source Water Assessments will be updated every five years to reflect any changes in the vulnerability and/or susceptibility of the drinking water source.


#### **REFERENCES**

 $A lask a \ Department \ of \ Community \ and \ Economic \ Development \ (ADCED), \ 2004 \ [WWW \ document]. \ URL \ http://www.dced.state.ak.us/cbd/commdb/CF\_COMDB.htm$ 

United States Environmental Protection Agency (EPA), 2004 [WWW document]. URL http://www.epa.gov/safewater/mcl.html.

# **APPENDIX A**

Kuparuk Operations Center Drinking Water Protection Area Location Map (Map 1)



# APPENDIX B

Contaminant Source Inventory and Risk Rankings (Tables 1 - 4)

# Contaminant Source Inventory for ConocoPhillips - Kuparuk OC

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Map Number	Comments
Pipelines (oil)	X28	X28-1	A	2	From operator submitted information.

# Contaminant Source Inventory and Risk Ranking for ConocoPhillips - Kuparuk OC Sources of Volatile Organic Chemicals

PWSID 330031.001

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Pipelines (oil)	X28	X28-1	A	Medium	2	From operator submitted information.

# Contaminant Source Inventory and Risk Ranking for ConocoPhillips - Kuparuk OC Sources of Heavy Metals, Cyanide and Other Inorganic Chemicals

PWSID 330031.001

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Pipelines (oil)	X28	X28-1	A	Low	2	From operator submitted information.


# Contaminant Source Inventory and Risk Ranking for ConocoPhillips - Kuparuk OC Sources of Other Organic Chemicals

PWSID 330031.001

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Pipelines (oil)	X28	X28-1	A	High	2	From operator submitted information.

### **APPENDIX C**

Kuparuk Operations Center
Drinking Water Protection Area
and Potential and Existing Contaminant Sources
(Map 2)



# APPENDIX D

# **Vulnerability Analysis and Contaminant Risks** (Charts 1-13)






Chart 2. Contaminant risks for ConocoPhillips, Kuparuk OC - Bacteria & Viruses Contaminant risks initially assumed to be low. What level of risk is associated Contaminant risks = with the highest and the next 0 pts + 0 pts highest risk source(s) of contaminants identified in Zone A (or floodplain)? Risk Rankings for Bacteria/Virus Contaminant Sources Identified Zone A Total Very Highs(s) 0 0 YES 0 High(s) Has there been two consecutive positive results Medium(s) 0 0 Increase susceptibility 50 for bacteria and viruses in Low(s) 0 0 pts recent sampling period(s)? VERY HIGH LOW **MEDIUM** HIGH 10 pts 20 pts 30 pts 40 pts Recent Bacteria Sampling ≥ 10 sources ≥ 20 sources ≥ 10 sources LOW Results + 10 pts + 5 pts + 5 pts ≥ 2 sources ≥ 5 sources ≥ 10 sources **MEDIUM** + 5 pts + 5 pts + 5 pts No recent sampling (inactive)  $\geq 1$  source ≥ 2 sources NO HIGH + 10 pts + 10 pts  $\geq 1$  source VERY HIGH + 10 pts Matrix Score 0 Note: Septic systems, sewerlines, and roads are each assigned a risk ranking for each individual contaminant source in the CSI. The VA, however, counts these contaminant sources as a group and assigns a calculated number of either "lows" or "mediums" based on the density.




Page 3 of 24

Chart 3. Vulnerability analysis for ConocoPhillips, Kuparuk OC - Bacteria & Viruses (Chart 1. Susceptibiltiy of the surface water source) Evaluate the susceptibility of the surface water Susceptibility of the surface source water Contaminant risks Vulnerability of drinking water surface water to contamination Susceptibility of surface water High 35 pts (Chart 2. Contaminant risks for surface water -Vulnerability of drinking water Bacteria & Viruses) surface water Evaluate 35 pts contaminant risks 35 Low Overall Vulnerability Ratings 80 to 100 pts very high Contaminant risks 60 to 79 pts high Low 40 to 59 pts medium 0 pts < 40 pts low



Page 5 of 24


Chart 4. Contaminant risks for ConocoPhillips, Kuparuk OC - Nitrates and Nitrites




Risk Levels for Nitrate/Nitrite Sources identified in Zones A and B								
	Zone A	Zone B	Total					
Very Highs(s)	0	0	0					
High(s)	0	0	0					
Medium(s)	0		0					
Low(s)	0		0					


	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	≥ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source + 10 pts


Matrix Score 0






Page 7 of 24






Page 9 of 24


Chart 6. Contaminant risks for ConocoPhillips, Kuparuk OC - Volatile Organic Chemicals




	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	≥ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source + 10 pts


Matrix Score 20





Page 11 of 24






Page 13 of 24

Chart 8. Contaminant risks for ConocoPhillips, Kuparuk OC - Heavy Metals, Cyanide and Other Inorganic Chemicals

+ 10 pts





0

	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	≥ 10 sources + 10 pts	≥ 10 sources 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source

N 6 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1.0
Matrix Score	10


Medium(s) Low(s)






Page 15 of 24

Chart 9. Vulnerability analysis for ConocoPhillips, Kuparuk OC - Heavy Metals, Cyanide and Other Inorganic Chemicals (Chart 1. Susceptibiltiy of the surface water source) Evaluate the susceptibility of the surface water Susceptibility of the surface source water Contaminant risks Vulnerability of drinking water surface water to contamination Susceptibility of surface water High 35 pts (Chart 8. Contaminant risks for surface water -Heavy Metals, Cyanide and Other Inorganic Vulnerability of drinking water Chemicals) surface water Evaluate 47 pts contaminant risks 45 Medium Overall Vulnerability Ratings 80 to 100 pts very high Contaminant risks 60 to 79 pts high Low 40 to 59 pts medium 12 pts < 40 pts low



Page 17 of 24


Chart 10. Contaminant risks for ConocoPhillips, Kuparuk OC - Synthetic Organic Chemicals




Risk Levels for SOC Sources identified in Zones A and C					
	Zone A	Zone B	Total		
Very Highs(s)	0	0	0		
High(s)	0	0	0		
Medium(s)	0	0	0		
Low(s)	0	0	0		


	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	≥ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source + 10 pts


Matrix Score 0






Page 19 of 24






Page 21 of 24


Chart 12. Contaminant risks for ConocoPhillips, Kuparuk OC - Other Organic Chemicals



	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	YERY HIGH 40 pts
LOW	≥ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source + 10 pts

Matrix Score 30





Page 23 of 24

