



Source Water Assessment

A Hydrogeologic Susceptibility and
Vulnerability Assessment for
Fort Wainwright Water Treatment Plant
Well 3405,
Fort Wainwright, Alaska

March 2004

PWSID 310918.8

DRINKING WATER PROTECTION PROGRAM REPORT Report 1289
Alaska Department of Environmental Conservation

Source Water Assessment for Fort Wainwright Water Treatment Plant Well 3405 Fort Wainwright, Alaska PWSID 310918.8

March 2004

DRINKING WATER PROTECTION PROGRAM REPORT Report 1289

The Drinking Water Protection Program (DWPP) is producing Source Water Assessments in compliance with the Safe Drinking Water Act Amendments of 1996. Each assessment includes a delineation of the source water area, an inventory of potential and existing contaminant sources that may impact the water, a risk ranking for each of these contaminants, and an evaluation of the potential vulnerability of these drinking water sources.

These assessments are intended to provide public water systems owners/operators, communities, and local governments with the best available information that may be used to protect the quality of their drinking water. The assessments combine information obtained from various sources, including the U.S. Environmental Protection Agency, Alaska Department of Environmental Conservation (ADEC), public water system owners/operators, and other public information sources. The results of this assessment are subject to change if additional data becomes available. It is anticipated this assessment will be updated every five years to reflect any changes in the vulnerability and/or susceptibility of public drinking water source. If you have any additional information that may affect the results of this assessment, please contact the Program Coordinator of DWPP, (907) 269-7521.

CONTENTS

Executive Summary Fort Wainwright Water Treatment Plant Public Drinking Water System Fort Wainwright Water Treatment Plant Protection Area		Page 1 1 1	Inventory of Potential and Existing Contaminant Sources Ranking of Contaminant Risks Vulnerability of Fort Wainwright Water Treatment Plant Drinking Water System References	Page 2 2 3 7			
			TAB	LES			
TABLE	1.	Definition of Zones		2			
 Susceptibility Contaminant Risks Overall Vulnerability 				4 4 4			
		Al	PPEN	DICES			
APPENDIX	A.	Fort Wainwright Water Trea	ıtment Pl	ant wells Drinking Water Protection Area (Map 1)			
		 B. Contaminant Source Inventory for Fort Wainwright Water Treatment Plant wells (T Contaminant Source Inventory and Risk Ranking for Fort Wainwright Water Treatm – Bacteria and Viruses (Table 2) Contaminant Source Inventory and Risk Ranking for Fort Wainwright Water Treatm – Nitrates/Nitrites (Table 3) Contaminant Source Inventory and Risk Ranking for Fort Wainwright Water Treatm – Volatile Organic Chemicals (Table 4) Contaminant Source Inventory and Risk Ranking for Fort Wainwright Water Treatm – Heavy Metals, Cyanide, and Other Inorganic Chemicals (Table 5) Contaminant Source Inventory and Risk Ranking for Fort Wainwright Water Treatm – Synthetic Organic Chemicals (Table 6) Contaminant Source Inventory and Risk Ranking for Fort Wainwright Water Treatm – Other Organic Chemicals (Table 7) C. Fort Wainwright Water Treatment Plant wells Drinking Water Protection Area and Pand Existing Contaminant Sources (Map 2) 					
	D.			ant Source Inventory and Risk Ranking for nt Plant wells Public Drinking Water Source			

Source Water Assessment for Fort Wainwright Water Treatment Plant Well 3405 Fort Wainwright, Alaska

Drinking Water Protection Program Alaska Department of Environmental Conservation

EXECUTIVE SUMMARY

This source water assessment provides an evaluation of the vulnerability to potential contamination of one of the nine wells making up the public water system serving Fort Wainwright Water Treatment Plant. This Class A (community) water system consists of nine wells in Fort Wainwright, Alaska. This report is an assessment of Well 3405 (PWSID 310918.8). Well 3405 is used primarily for fire protection but also serves as a backup well for the public water system. The well received a natural susceptibility rating of **Medium**. This rating is a combination of a susceptibility rating of Low for the actual wellhead and a Very High rating for the aquifer in which the well is drawing water from. Identified potential and current sources of contamination for this Fort Wainwright Water Treatment Plant well include: septic systems, residential areas, cement manufacturing, and DECrecognized contaminated sites. These are considered as sources of bacteria and viruses, nitrates and/or nitrites, volatile organic chemicals, heavy metals, cyanide and other inorganic chemicals, synthetic organic chemicals, and other organic chemicals. Combining the natural susceptibility of the well with the contaminant risk, the public water system for Fort Wainwright Water Treatment Plant wells received an overall vulnerability rating of **High** for bacteria and viruses, **Medium** for volatile organic chemicals, and Low for nitrates and/or nitrites, heavy metals, cyanide, and other inorganic chemicals, synthetic organic chemicals, and other organic chemicals.

FORT WAINWRIGHT WATER TREATMENT PLANT WELLS PUBLIC DRINKING WATER SYSTEM

Fort Wainwright Water Treatment Plant public water system is a Class A (community) water system. This is an assessment of one of the nine wells making up the public drinking water system serving Fort Wainwright. Well 3405 (PWSID 310918.8) is used primarily for fire protection but also serves as a backup well for the public water system. The well is located on Ile De France Avenue near its intersection with Santiago Avenue in Fort Wainwright, Alaska (T1S, R1E, Section 19) (See Map 1 of Appendix A). Fort Wainwright is located on the southeast side of the town of Fairbanks which is located in the Fairbanks North Star Borough

near the center of Alaska (Please see the inset of Map 1 in Appendix A for location). The Borough's current population is 82,840 making it the second-largest population center in the state (ADCED, 2002). Communities located within the Borough include: College, Eielson Air Force Base, Ester, Fairbanks, Fox, Harding Lake, Moose Creek, North Pole, Pleasant Valley, Salcha, and Two Rivers.

The Fairbanks area includes two distinct topographic areas: the alluvial plain between the Tanana River and the Chena River, and the uplands north of this alluvial plain. The Fort Wainwright Water Treatment Plant wells are located in the alluvial plain at an elevation of approximately 450 feet above sea level.

The depth of this well is unknown. Most wells in this area are screened in a combination of sand and gravel, and it is assumed these are also. The alluvial plain consists of alternating layers of silt, sand and gravel up to over 500 feet thick, in some locations overlain by 1 to 10 feet of silt or sandy silt or a few feet of peat (Glass and others, 1996). Discontinuous permafrost (perennially frozen areas) is also common in the alluvial plain. The depth to permafrost in these areas ranges between 2 and 45 feet below the ground surface with the thickness of the permafrost ranging between 5 and 265 feet (Pewe, T.L. 1958. Geology of the Fairbanks (D-2) Quadrangle, Alaska. USGS). Areas with discontinuous permafrost may locally affect the ground water flow directions.

Primarily the Tanana River, but also the Chena River contribute water to this alluvial aquifer. The Chena River typically only contributes water when its stage is high and the Tanana is low (Nelson, 1978). The Tanana River gets approximately 85% of its water from snowmelt of the Alaska Range and 15% from the Yukon-Tanana uplands (Anderson, 1970).

This public drinking water system serves approximately 6,000 residents through approximately 600 service connections.

FORT WAINWRIGHT WATER TREATMENT PLANT WELLS DRINKING WATER PROTECTION AREA

The pathways most likely for surface contamination to reach the groundwater are identified as the first step in determining a drinking water system's risk. These areas are determined by looking at the characteristics of the soil, groundwater, aquifer, and well.

The most probable area for contamination to reach the drinking water wells is the area that contributes water to the well, the groundwater capture zone. The groundwater capture zone is located in the area circling the well (the area influenced by pumping) and also the area of the water table upgradient of the well, usually forming a parabola shape.

There are many different ways of calculating the size of capture zones. This assessment uses a combination of two simple groundwater flow equations, the Thiem and uniform flow equations for all groundwater wells screened in unconsolidated material. The orientation of the capture zone is then drawn using a water table elevation map (if available) or a land surface elevation map of the area. The capture zone calculated in this assessment is only a best guess using the information and resources available to us, and may differ slightly from the actual capture zone.

The parameters used to calculate the shape of this capture zone are general for the whole alluvial plain and were obtained from various United State Geological Survey (USGS) reports, well logs in the area, and the Groundwater textbook by Freeze and Cherry (Freeze and Cherry, 1979).

The water table in the area of these Fort Wainwright Water Treatment Plant wells, the area between the Tanana and the Chena Rivers, is primarily influenced by the level of water flow in each river. The capture zones were drawn based on three separate configurations of the water table during various stages of the rivers: a period of high stage in the Chena River (October 14-17, 1986), high stage in the Tanana River (July 16-17, 1987), and low stages in both rivers (March 30-April 3, 1988) (Glass and others, 1996). High water levels in the Chena usually occur in the spring due to runoff from the uplands and in late summer due to rainstorms (Nelson, 1978). The Tanana usually experiences high flow during the hot, dry periods of mid-summer when maximum snowmelt from the Alaska Range occurs (Nelson, 1978). Groundwater in this area generally flows toward the northwest, from the Tanana River to the Chena River, however flow is reversed very near the Chena River during its high stage periods (Glass and others, 1996). These flow reversals are of short duration (i.e. days versus months) and of limited extent, generally within 1000 feet of the river (Nakanishi, et all, 1998).

Because of uncertainties and changing site conditions, a factor of safety is added to the groundwater capture zone to form the drinking water protection area for the well.

The protection areas established for wells are usually separated into four zones, limited by the watershed. These zones correspond to times-of-travel (TOT) of the water moving through the aquifer to the well (plus the factor of safety).

The following is a summary of the four zones for wells and the calculated time-of-travel for each:

Table 1. Definition of Zones

Zone	Definition
A	¹ / ₄ the distance for the 2-yr. time-of-travel
В	Less than 2 years time-of-travel
C	Less than 5 years time-of-travel
D	Less than 10 years time-of-travel

The time of travel for *contaminants* within the water varies with their unique physical and chemical characteristics.

The drinking water protection area outlined for the Fort Wainwright Water Treatment Plant wells on Map 1 of Appendix A will serve as the focus for voluntary protection efforts.

INVENTORY OF POTENTIAL AND EXISTING CONTAMINANT SOURCES

The Drinking Water Protection Program (DWPP) has completed an inventory of potential and existing sources of contamination within these Fort Wainwright Water Treatment Plant wells protection area. This inventory was completed through a search of agency records and other publicly available information. Potential drinking water contaminants are found within agricultural, residential, commercial, and industrial areas, but can also occur within areas that have little or no development.

For the basis of all Class A public water system assessments, six categories of drinking water contaminants were inventoried. They include:

- Bacteria and viruses;
- Nitrates and/or nitrites;
- Volatile organic chemicals;
- Heavy metals, cyanide, and other inorganic chemicals;
- Synthetic organic chemicals; and
- Other inorganic chemicals.

The sources are displayed on Map 2 of Appendix C and summarized in Table 1 of Appendix B.

RANKING OF CONTAMINANT RISKS

Once the potential and existing sources of contamination have been identified, they are each

assigned a ranking according to what type and level of risk they represent. Ranking of contaminant risks for a "potential" or "existing" source of contamination is a combination of toxicity and volume associated with that source. Rankings include:

Low;Medium;High; andVery High.

Bacteria and Viruses are only inventoried in Zones A and B because of their short life span. Only "Very High" and "High" rankings are inventoried within the outer Zone D due to the probability of contaminant dilution by the time the contaminants get to the well.

Tables 2 through 7 in Appendix B contain the ranking of inventoried potential and existing sources of contamination with respect to the six contaminant categories.

VULNERABILITY OF FORT WAINWRIGHT WATER TREATMENT PLANT WELLS DRINKING WATER SYSTEM

Vulnerability of a drinking water source to contamination is a combination of two factors:

- Natural susceptibility; and
- Contaminant risks.

Appendix D contains fourteen charts, which together form the 'Vulnerability Analysis' for a source water assessment for a public drinking water source. Chart 1 analyzes the 'Susceptibility of the Wellhead' to contamination by looking at the construction of the well and its surrounding area. Chart 2 analyzes the 'Susceptibility of the Aquifer' to contamination by looking at the properties of the aguifer and the presence of other wells or boreholes in the area. Chart 3 analyzes 'Contaminant Risks' for the drinking water source with respect to Bacteria and Viruses. The 'Contaminant Risks' portion of the analysis considers potential sources of contaminants as well as a review of the water system's contaminant sample results. Lastly, Chart 4 combines the results of the first three charts to produce the 'Vulnerability Analysis for Bacteria and Viruses'. Charts 5 through 14 contain the Contaminant Risks and Vulnerability Analyses for nitrates and nitrites, volatile organic chemicals, heavy metals, cyanide, and other inorganic chemicals, synthetic organic chemicals, and other organic chemicals, respectively.

A score for the Natural Susceptibility is reached by considering the properties of the well and the aquifer.

Susceptibility of the Wellhead (0 – 25 Points) (Chart 1 of Appendix D)

+

Susceptibility of the Aquifer (0-25 Points)(Chart 2 of Appendix D)

=

Natural Susceptibility (Susceptibility of the Well) (0-50 Points)

A ranking is assigned for the Natural Susceptibility according to the point score:

Natural Susceptibility Ratings								
40 to 50 pts	Very High							
30 to < 40 pts	High							
20 to < 30 pts	Medium							
< 20 pts	Low							

The wellhead for the Fort Wainwright Water Treatment Plant wells received a Low Susceptibility rating. The well is capped with a sanitary seal and the land surface is sloped away from the well, however the well is not grouted. A sanitary seal prevents potential contaminants from entering the well from the inside while sloping the land surface away from the well and grouting help to prevent potential contaminants from traveling down the outside of the well casing.

The aquifer in the area the Fort Wainwright Water Treatment Plant wells well is completed in received a Very High Susceptibility rating. The highly transmissive aquifer material (sand and gravel) in the area allows contaminants to travel downward from the surface with the precipitation and surface water runoff. The shallow water table allows potential contaminants to come into contact with the water table with little natural filtering where they can disperse quickly. Additionally, wells in the area can provide a quick pathway for contaminants to reach the aquifer. Table 2 summarizes the Susceptibility scores and ratings for Fort Wainwright Water Treatment Plant wells.

Table 2. Susceptibility

	Score	Rating
Susceptibility of the	5	Low
Wellhead		
Susceptibility of the	19	High
Aquifer		
Natural Susceptibility	24	Medium

The Contaminant Risk has been derived from an evaluation of the routine sampling results of the water system and the presence of potential sources of contamination. Contaminant risks to a drinking water source depend on the type and distribution of contaminant sources. Flow charts are used to assign a point score, and ratings are assigned in the same way as for the natural susceptibility:

Contaminant Risk Ratings								
40 to 50 pts	Very High							
30 to < 40 pts	High							
20 to < 30 pts	Medium							
< 20 pts	Low							

Table 3 summarizes the Contaminant Risks for each category of drinking water contaminants.

Table 3. Contaminant Risks

Category	Score	Rating
Bacteria and Viruses	50	Very High
Nitrates and/or Nitrites	10	Low
Volatile Organic Chemicals	18	Low
Heavy Metals, Cyanide, and		
Other Inorganic Chemicals	4	Low
Synthetic Organic Chemicals	4	Low
Other Organic Chemicals	4	Low

Finally, an overall vulnerability score is assigned for each water system by combining each of the contaminant risk scores with the natural susceptibility score:

Vulnerability of the Drinking Water Source to Contamination (0 - 100).

Again, rankings are assigned according to a point score:

Overall Vulnerability Ratings								
80 to 100 pts	Very High							
60 to < 80 pts	High							
40 to < 60 pts	Medium							
< 40 pts	Low							

Table 4 contains the overall vulnerability scores (0 – 100) and ratings for each of the six categories of drinking water contaminants. Note: scores are rounded off to the nearest five.

Table 4. Overall Vulnerability

Category	Score	Rating
Bacteria and Viruses	75	High
Nitrates and/or Nitrites	35	Low
Volatile Organic Chemicals	40	Medium
Heavy Metals, Cyanide, and		
Other Inorganic Chemicals	30	Low
Synthetic Organic Chemicals	30	Low
Other Organic Chemicals	30	Low

Bacteria and Viruses

Although there are no potential sources of Bacteria and Viruses identified in the protection area, coliforms (a bacteria) have been detected during recent sampling.

Only a small amount of bacteria and viruses are required to endanger public health. Coliforms are found naturally in the environment and although they aren't necessarily a health threat, it is an indicator of other potentially harmful bacteria in the water, more specifically, fecal coliforms and E. coli which only come from human and animal fecal waste (EPA, 2002). Harmful bacteria can cause diarrhea, cramps, nausea, headaches, or other symptoms (EPA, 2002). Routine sampling has most recently detected coliforms in the water on 9/4/01 (verified on 9/6/01). Fecal coliform and E. Coli were not detected in the water.

After combining the contaminant risk for bacteria and viruses with the natural susceptibility of the well, the overall vulnerability of the well to contamination is high.

Nitrates and Nitrites

The residential septic systems in Zone C represent the greatest risk of nitrates and nitrites for this source of public drinking water.

Nitrates are very mobile, moving at approximately the same rate as water. Nitrates have not been detected in significant concentrations during recent sampling

history for the Fort Wainwright Water Treatment Plant wells.

After combining the contaminant risk for nitrates and nitrites with the natural susceptibility of the well, the overall vulnerability of the well to contamination is low.

Volatile Organic Chemicals

The residential heating oil tanks represent the greatest risk for volatile organic chemical contamination to the well.

Volatile Organic Chemicals have not been detected during routine sampling of this water system. After combining the contaminant risk for volatile organic chemicals with the natural susceptibility of the well, the overall vulnerability of the well to contamination is medium.

Heavy Metals, Cyanide, and Other Inorganic Chemicals

The residential septic systems represent the greatest risk to heavy metals for this source of public drinking water.

Antimony and Beryllium have been detected in extremely small concentrations with respect to their Maximum Contaminant Levels (MCL). A MCL is the concentration of a contaminant allowed in the drinking water by the Environmental Protection Area (EPA). No other heavy metals were not detected during recent sampling.

After combining the contaminant risk for heavy metals, cyanide and other inorganic chemicals with the natural susceptibility of the well, the overall vulnerability of the well to contamination is low.

Synthetic Organic Chemicals

The septic systems and residential area represent the only identified risk to synthetic organic chemical contamination to this well.

Synthetic Organic Chemicals have not detected during routine sampling in the Fort Wainwright Water Treatment Plant water system.

After combining the contaminant risk for synthetic organic chemicals with the natural susceptibility of the well, the overall vulnerability of the well to contamination is low.

Other Organic Chemicals

The septic systems also represent the greatest risk for other organic chemical contamination to the well.

Other Organic Chemicals have not been detected during routine sampling in the Fort Wainwright Water Treatment Plant water system.

After combining the contaminant risk for other organic chemicals with the natural susceptibility of the well, the overall vulnerability of the well to contamination is low.

Existing Contaminated Sites


There are approximately 7 ADEC-recognized contaminated sites located within the protection area (displayed on Map 2 Appendix C). The locations of these sites are approximate and maybe at the center of the zipcode boundaries the actual site is located in. The ADEC Contaminated Sites program has identified, assessed, and is ensuring cleanup of these sites based on their prioritized order. Priority is based on risk to human health and the environment, including risk to public drinking water wells. Specific information on each site can be found on the internet at http://www.dec.state.ak.us/spar/cs/search/csites/csites_search.asp or by calling the ADEC Contaminated Sites Program at (907) 269-7658.

REFERENCES

- Alaska Department of Community and Economic Development (ADCED), 2002 [WWW document]. URL http://www.dced.state.ak,us/mra/CF_BLOCK.cfm.
- Anderson, G.S., 1970, Hydrologic reconnaissance of the Tanana basin, central Alaska: U.S. Geological Survey Hydrologic Investigations Atlas HA-319.
- Forbes, R.B. and Weber, F.R., 1981. Bedrock Geologic Map of the Fairbanks Mining District, Alaska. Funded by the State of Alaska, US Geological Survey, and The National Science Foundation.
- Freeze, R.A. and Cherry, J.A., 1979. Groundwater. Prentice-Hall, Englewood Cliffs, NJ.
- Glass, Roy L., Lilly, Micheal R., and Meyer, David F., 1996. Ground-Water Levels in an Alluvial Plain Between the Tanana and Chena Rivers Near Fairbanks, Alaska 1986-93. US Geological Survey Water Resources Investigations Report 96-4060, 39p.
- Nakanishi, Allan S. and Lilly, Micheal R., 1998. Estimate of Aquifer Properties by Numerically Simulating Ground-Water/Surface-Water Interactions, Fort Wainwright, Alaska. US Geological Survey Water Resources Investigations Report 98-4088, 27p.
- Nelson, Gordon L., 1978, Hydrologic Information for Land-Use Planning, Fairbanks Vicinity, Alaska. US Department of the Interior Geological Survey Open File Report 78-959, 47p.
- Pewe, T. L., 1958, Geologic map of the Fairbanks D-2 quadrangle, Alaska: U.S. Geol. Survey Geol. Quad. Map GQ-110, scale 1:63,360.
- United States Environmental Protection Agency (EPA), 2002 [WWW document]. URL http://www.epa.gov/safewater/mcl.html.

APPENDIX A

Fort Wainwright Water Treatment Plant wells Drinking Water Protection Area Location Map (Map 1)



APPENDIX B

Contaminant Source Inventory and Risk Ranking for Fort Wainwright Water Treatment Plant wells (Tables 1-7)

Table 1

Contaminant Source Inventory for Ft Wainwright / Wtr Trtmt Plt

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Map Number	Comments
Contaminated sites, DEC recognized, non-Superfund, non-RCRA	U04	U04-1	В	2	Off Luzon Avenue; FTWW (2P) Bldg. 3485 UST 321; File Number 108.26.020
Contaminated sites, DEC recognized, non-Superfund, non-RCRA	U04	U04-2	В	2	West of Luzon; FTWW (2P) Bldg. 3481 (RFA S-11); File Number 108.26.017
Contaminated sites, DEC recognized, non-Superfund, non-RCRA	U04	U04-3	В	2	Motor Pool; FTWW (2P) Bldg. 3483/3485; File Number 108.38.041
Contaminated sites, DEC recognized, non-Superfund, non-RCRA	U04	U04-4	В	2	East of Luzon Ave North of 3485; FTWW (2P) Bldg 3483; File Number 108.26.005
Residential Areas	R01	R01-1	C	2	Approximately 20 acres of residential area in Zone C
Septic systems (serves one single-family home)	R02		С	2	Estimated 3 residential septic systems based on number of parcels designated as residential
Tanks, heating oil, residential (above ground)	R08		С	2	Estimated 3 residential heating oil tanks based on number of parcels designated as residential
Contaminated sites, DEC recognized, non-Superfund, non-RCRA	U04	U04-5	C	2	1491 New Richardson Highway; Walsky Property; RecKey 1995310933501
Cement manufacturing	108	I08-1	D	2	900 Old Richardson Highway
Contaminated sites, DEC recognized, non-Superfund, non-RCRA	U04	U04-6	D	2	1792 Tootie Street; FDIC Tootie Street-North Pole; RecKey 1990310124801
Contaminated sites, DEC recognized, non-Superfund, non-RCRA	U04	U04-7	D	2	Mile 7.5 Old Richardson Hwy; Mat-Su Inc.; RecKey 1989310107301

Table 2

Contaminant Source Inventory and Risk Ranking for Ft Wainwright / Wtr Trtmt Plt Sources of Nitrates/Nitrites

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Septic systems (serves one single-family home)	R02		С	Low	2	Estimated 3 residential septic systems based on number of parcels designated as residential
Residential Areas	R01	R01-1	C	Low	2	Approximately 20 acres of residential area in Zone C

Contaminant Source Inventory and Risk Ranking for Ft Wainwright / Wtr Trtmt Plt Sources of Volatile Organic Chemicals

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Septic systems (serves one single-family home)	R02		С	Low	2	Estimated 3 residential septic systems based on number of parcels designated as residential
Tanks, heating oil, residential (above ground)	R08		C	Medium	2	Estimated 3 residential heating oil tanks based on number of parcels designated as residential
Residential Areas	R01	R01-1	C	Low	2	Approximately 20 acres of residential area in Zone C
Cement manufacturing	108	I08-1	D	High	2	900 Old Richardson Highway

Table 4

Contaminant Source Inventory and Risk Ranking for Ft Wainwright / Wtr Trtmt Plt Sources of Heavy Metals, Cyanide and Other Inorganic Chemicals

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Septic systems (serves one single-family home)	R02		С	Low	2	Estimated 3 residential septic systems based on number of parcels designated as residential
Residential Areas	R01	R01-1	С	Low	2	Approximately 20 acres of residential area in Zone C

Table 5

Contaminant Source Inventory and Risk Ranking for Ft Wainwright / Wtr Trtmt Plt Sources of Synthetic Organic Chemicals


Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Septic systems (serves one single-family home)	R02		С	Low	2	Estimated 3 residential septic systems based on number of parcels designated as residential
Residential Areas	R01	R01-1	С	Low	2	Approximately 20 acres of residential area in Zone C

Contaminant Source Inventory and Risk Ranking for Ft Wainwright / Wtr Trtmt Plt Sources of Other Organic Chemicals

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Septic systems (serves one single-family home)	R02		С	Low	2	Estimated 3 residential septic systems based on number of parcels designated as residential
Residential Areas	R01	R01-1	C	Low	2	Approximately 20 acres of residential area in Zone C
Cement manufacturing	108	I08-1	D	Very High	2	900 Old Richardson Highway

APPENDIX C

Fort Wainwright Water Treatment Plant wells
Drinking Water Protection Area
and Potential and Existing Contaminant Sources
(Map 2)



APPENDIX D

Vulnerability Analysis for Fort Wainwright Water Treatment Plant wells Public Drinking Water Source (Charts 1-14)

Chart 1. Susceptibility of the wellhead - Ft. Wainwright 3405




Chart 2. Susceptibility of the aquifer - Ft. Wainwright 3405








Chart 3. Contaminant risks for Ft. Wainwright 3405 - Bacteria & Viruses






Page 4 of 25





Page 6 of 25


Chart 5. Contaminant risks for Ft. Wainwright 3405 - Nitrates and Nitrites





sk Levels for Contaminant Sources identified in Zones A, B and C					
	Zone A	Zones B&C	Total		
Very Highs(s)	0	0	0		
High(s)	0	0	0		
Medium(s)	0	1	1		
Low(s)	0	2	2		


	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	≥ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source + 10 pts

Matrix Score	20
--------------	----










Page 10 of 25


Chart 7. Contaminant risks for Ft. Wainwright 3405 - Volatile Organic Chemicals




sk Levels for Contaminant Sources identified in Zones A, B and C					
	Zone A	Zones B&C	Total		
Very Highs(s)	0	0	0		
High(s)	0	0	0		
Medium(s)	0	3	3		
Low(s)	0	2	2		


	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	≥ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source + 10 pts


Matrix Score	25
--------------	----






Page 12 of 25





Page 14 of 25


Chart 9. Contaminant risks for Ft. Wainwright 3405 - Heavy Metals, Cyanide and Other Inorganic Chemicals





isk Levels for Contaminant Sources identified in Zones A, B and C					
	Zone A	Zones B&C	Total		
Very Highs(s)	0	0	0		
High(s)	0	0	0		
Medium(s)	0	0	0		
Low(s)	0	2	2		


	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	≥ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source + 10 pts

Matrix Score 10










Page 18 of 25


Chart 11. Contaminant risks for Ft. Wainwright 3405 - Synthetic Organic Chemicals





sk Levels for Contaminant Sources identified in Zones A, B and C					
	Zone A	Zones B&C	Total		
Very Highs(s)	0	0	0		
High(s)	0	0	0		
Medium(s)	0	0	0		
Low(s)	0	2	2		


	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	≥ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source + 10 pts

Matrix Score	10
--------------	----










Page 22 of 25

Chart 13. Contaminant risks for Ft. Wainwright 3405 - Other Organic Chemicals



Risk Levels for Contaminant Sources identified in Zones A, B and C					
	Zone A	Zones B&C	Total		
Very Highs(s)	0	0	0		
High(s)	0	0	0		
Medium(s)	0	0	0		
Low(s)	0	2	2		

	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	≥ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source + 10 pts

Matrix Score	10
--------------	----

