



Source Water Assessment

A Hydrogeologic Susceptibility and
Vulnerability Assessment for
Eaglecrest Ski Area,
Juneau, Alaska
PWSID #110643

DRINKING WATER PROTECTION PROGRAM REPORT NO. 715

Alaska Department of Environmental Conservation

Source Water Assessment for Eaglecrest Ski Area Juneau, Alaska PWSID #110643

DRINKING WATER PROTECTION PROGRAM REPORT NO. 715

The Drinking Water Protection Program (DWPP) is producing Source Water Assessments in compliance with the Safe Drinking Water Act Amendments of 1996. Each assessment includes a delineation of the source water area, an inventory of potential and existing contaminant sources that may impact the water, a risk ranking for each of these contaminants, and an evaluation of the potential vulnerability of these drinking water sources.

These assessments are intended to provide public water systems owners/operators, communities, and local governments with the best available information that may be used to protect the quality of their drinking water. The assessments combine information obtained from various sources, including the U.S. Environmental Protection Agency, Alaska Department of Environmental Conservation (ADEC), public water system owners/operators, and other public information sources. The results of this assessment are subject to change if additional data becomes available. It is anticipated this assessment will be updated every five years to reflect any changes in the vulnerability and/or susceptibility of public drinking water source. If you have any additional information that may affect the results of this assessment, please contact the Program Coordinator of DWPP, (907) 269-7521.

CONTENTS

	Page
SECTION	
Executive Summary	1
Eaglecrest Ski Area Public Drinking Water System	1
Eaglecrest Ski Area Drinking Water Protection Area	1
Inventory of Potential and Existing Contaminant Sources	2
Ranking of Contaminant Risks	2
Vulnerability of Eaglecrest Ski Area Drinking Water System	2
References	4

TABLES

TABLE	1. Definition of Zones	2
	2. Susceptibility	2
	3. Contaminant Risks	2
	4. Overall Vulnerability	3

APPENDICES

APPENDIX	A. Eaglecrest Ski Area Drinking Water Protection Area (Map 1)	
	B. Vulnerability Analysis for Contaminant Source Inventory and Risk Ranking for Eaglecrest Ski Area Public Drinking Water Source (Charts 1 – 7)	

Source Water Assessment for Eaglecrest Ski Area, Juneau, Alaska

Drinking Water Protection Program Alaska Department of Environmental Conservation

EXECUTIVE SUMMARY

The public water system for Eaglecrest Ski Area is a Class B (transient/non-community) water system consisting of one surface water intake from Eagle Creek located in the middle of Douglas Island across the Gastineau Channel from Juneau, Alaska. The surface water intake received a susceptibility rating of **Very High**. A rating of high to Very High is typical for all surface water systems. No potential and current sources of contaminants for Eaglecrest Ski Area public drinking water source were identified. However, identified potential and existing sources of contamination may include sources of bacteria and viruses, nitrates and/or nitrites, and volatile organic chemicals. Contaminant sources could potentially contribute bacteria and viruses, nitrates and/or nitrites, and volatile organic chemicals into the surface water source. Overall, the public water sources for Eaglecrest Ski Area received a vulnerability rating of **Medium** for bacteria and viruses; nitrates and nitrites; and volatile organic chemicals.

EAGLECREST SKI AREA PUBLIC DRINKING WATER SYSTEM

Eaglecrest Ski Area public water system is a Class B (transient/non-community) water system. The system consists of one surface water intake located on Douglas Island, across from Juneau, Alaska (See Map 1 of Appendix A).

Downtown Juneau averages about 92 inches of precipitation per year and approximately 101 inches of snow. The groundwater sources underlying the area are recharged through the infiltration of precipitation and surface water. Groundwater sources in the region generally occur in the fractured bedrock and unconsolidated sediments deposited by glaciers and/or rivers. Douglas Island topography varies from near sea level along the Gastineau Channel to approximately 3,100 feet to mountains surrounding the lodge.

There is no Sanitary Survey available for this surface water intake. However, as-builts for water system improvements indicate that the intake was adequately constructed. An adequately constructed intake may provide protection against debris and contaminants

from entering the system. The raw water is treated by filtration and chlorination. There is a potential for runoff within the area surrounding the surface water.

This system operates seasonally from October through April.

EAGLECREST SKI AREA DRINKING WATER PROTECTION AREA

In order to evaluate whether a drinking water source is at risk, we must first evaluate what are the most likely pathways for surface contamination to reach the creek. These areas are determined by looking at the characteristics of the creek, surrounding contaminant sources, and the intake.

The most probable area for contamination to reach the drinking water system is the area that contributes water to the surface water body that water is being drawn from. This area is designated as the Drinking Water Protection Area (DWPA). Because releases of contaminants within the DWPA are most likely to impact the drinking water system, this area will serve as the focus for voluntary protection efforts.

The size and shape of the DWPAs were established based on aerial distances from the surface water body, and the watershed that recharges the surface water body. Please refer to the Guidance Manual for Class B Public Water Systems for additional information.

The DWPAs established for surface water systems by the ADEC are separated into three zones. These zones correspond to different distances from the surface water body, and the entire watershed that recharges the surface water body. The following is a summary of the three DWPA zones and their definitions.

Table 1. Definition of Zones

Zone	Definition
A	1,000 feet from the Surface Water Body
B	1 mile from the Surface Water Body
C	Entire Watershed

The DWPA for Eaglecrest Ski Area extends over the entire watershed. Development in the vicinity of the surface water intake is downstream of the intake (See Map 1 of Appendix A).

INVENTORY OF POTENTIAL AND EXISTING CONTAMINANT SOURCES

The Drinking Water Protection Program has completed an inventory of potential and existing sources of contamination within the Eaglecrest Ski Area DWPA. This inventory was completed through a search of agency records and other publicly-available information. Potential sources of contamination to the drinking water source include a wide range of categories and types. Potential drinking water contaminants are found within agricultural, residential, commercial, and industrial areas, but can also occur within areas that have little or no development.

For the basis of all Class B public water system assessments, three categories of drinking water contaminants were inventoried. They include:

- Bacteria and viruses;
- Nitrates and/or nitrites;
- Volatile organic chemicals

RANKING OF CONTAMINANT RISKS

Once the potential and existing sources of contamination have been identified, they are assigned a ranking according to what type and level of risk they represent. Ranking of contaminant risks for a “potential” or “existing” source of contamination is a function of toxicity and volumes of specific contaminants associated with that source. Rankings include:

- Low;
- Medium;
- High; and
- Very High.

The TOT for contaminants within the water varies and is dependent on the physical and chemical characteristics of each contaminant. Bacteria and Viruses are only inventoried in Zones A and B because of their short life span.

VULNERABILITY OF EAGLECREST SKI AREA DRINKING WATER SYSTEM

Vulnerability of a drinking water source to contamination is a combination of two factors:

- Natural susceptibility; and
- Contaminant risks.

Each of the three categories of drinking water contaminants has been analyzed and an overall vulnerability score of 30 to 100 is ultimately assigned:

Natural Susceptibility (30 – 50 points)

+

Contaminant Risks (0 – 50 points)

=

Vulnerability of the Drinking Water Source to Contamination (30 – 100).

A score for the Natural Susceptibility is achieved by analyzing the properties of the surface water source.

Natural Susceptibility
(Susceptibility of the Surface Water Source)
(30 – 50 Points)

The surface water intake for Eaglecrest Ski Area is in Eagle Creek. Because Eagle Creek is recharged by surface water runoff and precipitation, contaminants at or near the creek have the potential to adversely impact this drinking water source. Table 2 shows the Overall Susceptibility score and rating for Eaglecrest Ski Area.

Table 2. Natural Susceptibility

	Score	Rating
Natural Susceptibility	45	Very High

Contaminant risks to a drinking water source depend on the type, number or density, and distribution of contaminant sources. This data has been derived from an examination of existing or historical contamination that has been detected at the drinking water source through routine sampling. It also evaluates potential sources of contamination. Table 3 summarizes the Contaminant Risks for each category of drinking water contaminants.

Table 3. Contaminant Risks

Category	Score	Rating
Bacteria and Viruses	0	Low
Nitrates and/or Nitrites	1	Low
Volatile Organic Chemicals	0	Low

Appendix D contains seven charts, which together form the ‘Vulnerability Analysis’ for a source water assessment for a public drinking water source. Chart 1 analyzes the ‘Susceptibility of the Surface Water Source’ to contamination by looking at the construction of the intake and its surrounding area and naturally-occurring attributes of the water source and influences on the groundwater system that might lead to contamination. Chart 2 analyzes ‘Contaminant Risks’

for the drinking water source with respect to bacteria and viruses. The ‘Contaminant Risks’ portion of the analysis considers potential sources of contaminants as well as a review of contamination that has or may have occurred, but has not arrived or been detected at the intake. Chart 3 contains the ‘Vulnerability Analysis for Bacteria and Viruses.’ Charts 4 through 7 contain the Contaminant Risks and Vulnerability Analyses for nitrates and nitrites and volatile organic chemicals, respectively.

Table 4 contains the overall vulnerability scores (30 – 100) and ratings for each of the three categories of drinking water contaminants. Note: scores are rounded off to the nearest five.

Table 4. Overall Vulnerability

Category	Score	Rating
Bacteria and Viruses	45	Medium
Nitrates and Nitrites	45	Medium
Volatile Organic Chemicals	45	Medium

Bacteria and Viruses

The contaminant risk for bacteria and viruses is **Low** with no unnatural sources of contamination identified (See Chart 2 – Contaminant Risks for Bacteria and Viruses in Appendix D).

Only a small amount of bacteria and viruses are required to endanger public health. Bacteria and viruses have not been detected during recent water sampling of the system at the Eaglecrest Ski Area. Combining the contaminant risks with the overall natural susceptibility of the surface water source, the vulnerability of the surface water source to contamination by bacteria and viruses is **Medium**.

Nitrates and Nitrites

The contaminant risk for nitrates and nitrites is **Low** with no unnatural sources of contamination identified (See Chart 4 - Contaminant Risks for Nitrates and/or Nitrites in Appendix D).

Sampling history for Eaglecrest Ski Area indicates that nitrates have been detected in the water, but only in very low concentrations (most recently at 0.160 mg/L on 12/13/1999) or 2% of the Maximum Contaminant Level (MCL). The MCL is the maximum level of contaminant that is allowed to exist in drinking water and still be consumed by humans without harmful health effects. Due to the high solubility and weak retention by soil, nitrates are very mobile, moving at approximately the same rate as water.

After combining the contaminant risk for nitrates and nitrites with the natural susceptibility of the surface water source, the overall vulnerability of the surface water source to contamination by nitrates and nitrites is **Medium**.

Volatile Organic Chemicals

The contaminant risk for volatile organic chemicals is **Low** with no sources of contamination identified (See Chart 6 – Contaminant Risks for Volatile Organic Chemicals in Appendix D).

Sampling history indicates that volatile organic chemicals have not been detected in the water. Combining the contaminant risk for volatile organic chemicals with the natural susceptibility of the surface water source, the overall vulnerability of the surface water source to contamination by volatile organic chemicals is **Medium**.

REFERENCES

Alaska Department of Community and Economic Development (ADCED), 2002 [WWW document]. URL http://www.dced.state.ak.us/mra/CF_BLOCK.cfm.

Alaska Geospatial Data Clearinghouse, 2003. URL: <http://agdc.usgs.gov/data/datasets.html>.



Gehrels, G.E., Berg, H.C., Geologic Map of Southeastern Alaska: U.S. Geological Survey Map (scale 1:600,000), Map I-1867, 1sheet.

King, P.B., compiler, 1969, Tectonic map of North America: US Geological Survey Map, (scale 1:5,000,000) 2 sheets.

United States Environmental Protection Agency (EPA), 2002 [WWW document]. URL: <http://www.epa.gov/safewater/mcl.html>.

APPENDIX A

Eaglecrest Ski Area Drinking Water Protection Area Location Map (Map 1)



Map 1: Eaglecrest Ski Area Drinking Water Protection Areas

PWSID: 110643.001




1:7,098

Data Sources:
Background image - USGS 1:25,000 mapping

Protection zones were delineated based upon streams noted on USGS 1:25,000 mapping.
For this PWS, Zone C (the entire watershed) covers the same area as Zone B (areas within 1 Mile of the stream).

- Legend**
- Public Drinking Water Systems
 - Zone A Protection Area**
1000 Feet from Surface Water
 - Zone B Protection Area**
1 Mile from Surface Water Body
 - Zone C Protection Area**
Entire Watershed



May 7, 2003

APPENDIX B

Vulnerability Analysis for Eaglecrest Ski Area Public Drinking Water Source (Charts 1-7)

Chart 1. Susceptibility of the surface water source - Eaglecrest Ski Area







Chart 2. Contaminant risks for Eaglecrest Ski Area - Bacteria & Viruses





Risk Rankings for Contaminant Sources Identified in Zones A and B			
	Zone A	Zone B	Total
Very High(s)	0	0	0
High(s)	0	0	0
Medium(s)	0	0	0
Low(s)	0	0	0

	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	³ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	----
MEDIUM	----	≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH	----	----	≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH	----	----	----	≥ 1 source + 10 pts


Matrix Score 0

Note: Septic systems, sewerlines, and roads are each assigned a risk ranking for each individual contaminant source in the CSI. The VA, however, counts these contaminant sources as a group and assigns a calculated number of either "lows" or "mediums" based on the density.

Chart 2. Contaminant risks for Eaglecrest Ski Area - Bacteria & Viruses



Contaminant Risk Ratings	
40 to 50 pts	very high
30 to < 40 pts	high
20 to < 30 pts	medium
< 20 pts	low



Existing	0 pts
+	
Potential	0 pts
=	
Contaminant Risk	0 pts

Chart 3. Vulnerability analysis for Eaglecrest Ski Area - Bacteria & Viruses




Chart 4. Contaminant risks for Eaglecrest Ski Area - Nitrates and Nitrites




Chart 4. Contaminant risks for Eaglecrest Ski Area - Nitrates and Nitrites






Chart 4. Contaminant risks for Eaglecrest Ski Area - Nitrates and Nitrites




Chart 5. Vulnerability analysis for Eaglecrest Ski Area - Nitrates and Nitrites




Chart 6. Contaminant risks for Eaglecrest Ski Area - Volatile Organic Chemicals





Chart 6. Contaminant risks for Eaglecrest Ski Area - Volatile Organic Chemicals



Risk Levels for Contaminant Sources identified in Zones A and B			
	Zone A	Zone B	Total
Very High(s)	0	0	0
High(s)	0	0	0
Medium(s)	0	0	0
Low(s)	0	0	0

	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	³ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	----
MEDIUM	----	≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH	----	----	≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH	----	----	----	≥ 1 source + 10 pts

Matrix Score 0

Note: Septic systems, sewerline, and roads are each assigned a risk ranking for each individual contaminant source in the CSI. The VA, however, counts these contaminant sources as a group and assigns a calculated number of either "lows" or "mediums" based on the density.





Chart 6. Contaminant risks for Eaglecrest Ski Area - Volatile Organic Chemicals





Chart 7. Vulnerability analysis for Eaglecrest Ski Area - Volatile Organic Chemicals

