

Source Water Assessment

A Hydrogeologic Susceptibility and Vulnerability Assessment for Stoney Creek R.V. Park Public Drinking Water System, Seward Area, Alaska PWSID # 249049.001

DRINKING WATER PROTECTION REPORT 1716

Alaska Department of Environmental Conservation

December, 2008

Source Water Assessment for Stoney Creek R.V. Park Public Drinking Water System Seward Area, Alaska PWSID# 249049.001

DRINKING WATER PROTECTION REPORT 1716

The Drinking Water Protection (DWP) section of the Drinking Water Program is producing Source Water Assessments in compliance with the Safe Drinking Water Act Amendments of 1996. Each assessment includes a delineation of the source water area, an inventory of potential and existing contaminant sources that may impact the water, a risk ranking for each of these contaminants, and an evaluation of the potential vulnerability of these drinking water sources.

These assessments are intended to provide public water systems owners/operators, communities, and local governments with the best available information that may be used to protect the quality of their drinking water. The assessments combine information obtained from various sources, including the U.S. Environmental Protection Agency, Alaska Department of Environmental Conservation (DEC), public water system owners/operators, and other public information sources. The results of this assessment are subject to change if additional data becomes available. It is anticipated this assessment will be updated every five years to reflect any changes in the vulnerability and/or susceptibility of public drinking water source. If you have any additional information that may affect the results of this assessment, please contact the DWP staff at the following number: 1-866-956-7656.

December, 2008

CONTENTS

Page

Executive Summary	Ranking of Contaminant Risks2 Vulnerability of Stoney Creek R.V. Park Drinking
Stoney Creek R.V. Park Drinking Water Protection	Water System2
Area	References
Inventory of Potential and Existing Contaminant Sources	Appendix B
5041005	Appendix C11

TABLES

2
3
3
3
•

APPENDICES

APPENDIX

- A. Stoney Creek R.V. Park Drinking Water Protection Area (Map A)
- B. Contaminant Source Inventory for Stoney Creek R.V. Park (Table 1) Contaminant Source Inventory and Risk Ranking for Stoney Creek R.V. Park – Bacteria and Viruses (Table 2) Contaminant Source Inventory and Risk Ranking for Stoney Creek R.V. Park – Nitrates/Nitrites (Table 3) Contaminant Source Inventory and Risk Ranking for Stoney Creek R.V. Park – Volatile Organic Chemicals (Table 4)
- C. Stoney Creek R.V. Park Drinking Water Protection Area and Potential and Existing Contaminant Sources (Map C)

Source Water Assessment for Stoney Creek R.V. Park Source of Public Drinking Water, Seward Area, Alaska

Drinking Water Protection Alaska Department of Environmental Conservation

EXECUTIVE SUMMARY

The public water system for Stoney Creek R.V. Park is a Class B (transient/non-community) water system consisting of one well located on the Seward Highway about 4 miles north of Seward, Alaska. The wellhead received a susceptibility rating of Low and the aquifer received a susceptibility rating of Very High. Combining these two ratings produces a Medium rating for the natural susceptibility of the well. Identified potential and current sources of contaminants for Stoney Creek R.V. Park public drinking water source include: inferred septic systems, assumed heating oil tanks, roads, and a campground/R.V. park. These identified potential and existing sources of contamination are considered as sources of bacteria and viruses, nitrates and/or nitrites, and volatile organic chemicals. Overall, the public water sources for Stoney Creek R.V. Park received a vulnerability rating of Low for bacteria and viruses, Medium for nitrates and nitrites, and Medium for volatile organic chemicals. This assessment of contaminant risks can be used as a foundation for local voluntary protection efforts as well as a basis for the continuous efforts on the part of Stoney Creek R.V. Park to protect public health.

STONEY CREEK R.V. PARK PUBLIC DRINKING WATER SYSTEM

The Stoney Creek R.V. Park public water system is a Class B (transient/non-community) water system. The system consists of one well located located on the Seward Highway, about 4 miles north of Seward, Alaska (see Map A in Appendix A). Seward is located at the head of Resurrection Bay in southcentral Alaska and is part of the Kenai Peninsula Borough. The city lies 125 road miles south of Anchorage and has a population of 2,661, while the Borough has a population of 52,407. Average annual precipitation in the area is 66 inches and 80 inches of snow. Temperatures range from 17 to 38 degrees in the winter and from 49 to 63 degrees in the summer (ADCCED, 2008).

Water is supplied within Seward by nine wells. Sewage is piped to a secondary treatment lagoon. Outside of Seward, individual wells and septic systems are common. A refuse transfer station is located on Hemlock Street. Electricity is provided by Seward Electric System, which buys it from Chugach Electric (ADCCED, 2008).

According to the well log, the well extends approximately 126 feet below the ground surface and is completed in a semi-confined aquifer. The sanitary survey (09/08/2004) for this system indicates that a sanitary seal is installed, the land surface is appropriately sloped away from the well, and the well is grouted according to DEC regulations.

This system operates from May to September and serves 2 residents and up to 240 non-residents through 82 service connections.

STONEY CREEK R.V. PARK DRINKING WATER PROTECTION AREA

In order to evaluate whether a drinking water source is at risk, we must first evaluate what are the most likely pathways for surface contamination to reach the groundwater. These areas are determined by looking at the characteristics of the soil, groundwater, aquifer, and well.

The most probable area for contamination to reach the drinking water well is the drinking water protection area. The drinking water protection area is the area circling the well (the area influenced by pumping) and also the area upgradient of the well, usually forming a parabola shape. Because releases of contaminants within the protection area are most likely to impact the well, this area will serve as the focus for voluntary protection efforts.

There are many different methods for calculating the size of protection areas. Drinking Water Protection (DWP) uses a combination of two simple groundwater flow equations, the Thiem and uniform flow equations for all groundwater wells screened in unconsolidated material. The orientation of the protection zone is then drawn using a water table elevation map (if available) or a land surface elevation map of the area. The protection zone calculated by the DWP is an estimate using the available information and resources, and may differ slightly from the actual capture zone. Because of uncertainties and changing site conditions, a factor of safety is added to the protection zone to form the drinking water protection area for the well.

The parameters used to calculate the shape of this protection zone are general for the whole alluvial plain and were obtained from various United States Geological Survey (USGS) reports, area well logs, and the Groundwater textbook by Freeze and Cherry (Freeze and Cherry, 1979).

The protection areas established for wells by the DEC are usually separated into two zones, limited by the watershed. These zones correspond to differences in the time-of-travel (TOT) of the water moving through the aquifer to the well. An analytical calculation was used to determine the size and shape of the protection area.

The time-of-travel for contaminants within the water varies and is dependent on the physical and chemical characteristics of each contaminant. The following is a summary of the two protection area zones for wells and the calculated time-of-travel for each:

Table 1. Definition of Zones

Zone	Definition
А	Several months time-of-travel
В	Less than the 2 year time-of-travel

The drinking water protection area for Stoney Creek R.V. Park was determined using an analytical calculation and includes Zones A and B (See Map A of Appendix A).

INVENTORY OF POTENTIAL AND EXISTING CONTAMINANT SOURCES

Drinking Water Protection has completed an inventory of potential and existing sources of contamination within the Stoney Creek R.V. Park drinking water protection area. This inventory was completed through a search of agency records and other publicly available information. Potential sources of contamination to the drinking water aquifer include a wide range of categories and types. Potential drinking water contaminants are found within agricultural, residential, commercial, and industrial areas, but can also occur within areas that have little or no development.

For the basis of all Class B public water system assessments, the following three categories of drinking water contaminants were inventoried:

- Bacteria and viruses;
- Nitrates and/or nitrites;
- Volatile organic chemicals

The sources are displayed on Map C of Appendix C and summarized in Table 1 of Appendix B.

RANKING OF CONTAMINANT RISKS

Once the potential and existing sources of contamination have been identified, they are assigned a ranking according to what type and level of risk they represent. Ranking of contaminant risks for a "potential" or "existing" source of contamination is a function of toxicity and volumes of specific contaminants associated with that source. Rankings include:

- Low;
- Medium;
- High; and
- Very High.

Tables 2 through 4 in Appendix B contain the ranking of potential and existing sources of contamination with respect to bacteria and viruses, nitrates and/or nitrites, and volatile organic chemicals.

VULNERABILITY OF STONEY CREEK R.V. PARK DRINKING WATER SYSTEM

Vulnerability of a drinking water source to contamination is a combination of two factors:

- Natural Susceptibility; and
- Contaminant Risks.

A score for the Natural Susceptibility of the well is reached by considering the properties of the well and the aquifer.

> Susceptibility of the Wellhead (0-25 Points) + Susceptibility of the Aquifer (0-25 Points) =

Natural Susceptibility of the Well (0-50 Points)

A ranking is assigned for the Natural Susceptibility according to the point score:

Natural Susceptibility Ratings					
40-50 pts	Very High				
30 to < 40 pts	High				
20 to < 30 pts	Medium				
< 20 pts	Low				

Factors contributing to the susceptibility of the wellhead are: whether the sanitary seal is in place, protection from flooding, and if the well casing is properly grouted.

The wellhead for the Stoney Creek R.V. Park received a **Low** susceptibility rating. The sanitary survey indicates that a sanitary seal is installed, the land surface is appropriately sloped away from the well, and the well is grouted according to DEC regulations. Sanitary seals prevent potential contaminants from entering the well, while sloping of the land surface away from the wellhead provides adequate surface water drainage, and concrete or grouting around the wellhead helps to prevent potential contaminants from traveling down the outside of the well casing.

Factors contributing to the susceptibility of the aquifer are: whether the aquifer is confined or unconfined, whether the well is completed in unconsolidated or fractured bedrock, whether wells and bore holes are penetrating the aquifer and, if applicable, the confining layer.

The aquifer that the Stoney Creek R.V. Park system is in is unconfined. It received a **Very High** susceptibility rating because of its semi-confined nature and the presence of other wells penetrating the vadose zone within the protection area. Because a semi-confined aquifer is partially recharged by surface water and precipitation that migrates downward from the surface, it is susceptible to contamination from outside sources. Furthermore, the presence of other wells penetrating the vadose zone of the protection area can allow contaminants to travel into the shared aquifer with precipitation and runoff.

Table 2 summarizes the Susceptibility scores and ratings for the Stoney Creek R.V. Park system.

Table 2. Susceptibility

	Score	Rating
Susceptibility of the	0	Low
Wellhead		
Susceptibility of the	25	Very High
Aquifer		
Natural Susceptibility	25	Medium

Contaminant risks are derived from an evaluation of the routine sampling results of the water system and the presence of potential sources of contamination. Contaminant risks to a drinking water source depend on the type and distribution of contaminant sources. Flow charts are used to assign a point score, and ratings are assigned in the same way as for the natural susceptibility:

Contaminant Risk Ratings					
40-50 pts	Very High				
30 to < 40 pts	High				
20 to < 30 pts	Medium				
< 20 pts	Low				

Table 3 summarizes the Contaminant Risks for each category of drinking water contaminants for the Stoney Creek R.V. Park system.

Table 3. Contaminant Risks

Category	Score	Rating
Bacteria and Viruses	12	Low
Nitrates and/or Nitrites	15	Low
Volatile Organic Chemicals	25	Medium

Finally, an overall vulnerability score is assigned for each water system by combining each of the contaminant risk scores with the natural susceptibility score:

Natural Susceptibility (0-50 Points)

Vulnerability of the Drinking Water Source to Contamination (0-100 Points)

Again, rankings are assigned according to a point score:

0	verall Vulnerab	ility Ratings
80-	-100 pts	Very High
60	to < 80 pts	High
40	to < 60 pts	Medium
< 4	0 pts	Low

Table 4 contains the overall vulnerability scores (0-100) and ratings for each of the three categories of drinking water contaminants for the Stoney Creek R.V. Park system. Note: scores are rounded off to the nearest five.

Table 4. Overall Vulnerability

Category	Score	Rating
Bacteria and Viruses	35	Low
Nitrates and/or Nitrites	40	Medium
Volatile Organic Chemicals	50	Medium

Bacteria and Viruses

The contaminant risk for bacteria and viruses is **Low** with the septic systems, roads, and campground contributing to the risk to the drinking water well.

Coliforms (a bacteria) are found naturally in the environment and although they aren't necessarily a health threat, they are an indicator of other potentially harmful bacteria in the water, more specifically, fecal coliforms and E. coli, which only come from human and animal fecal waste. Harmful bacteria can cause diarrhea, cramps, nausea, headaches, or other symptoms (EPA, 2008).

Only a small amount of bacteria and viruses are required to endanger public health. Positive samples increase the overall vulnerability of the drinking water source, indicating that the source is susceptible to bacteria and virus contamination. Bacteria and viruses have not been detected during recent water sampling of the system at Stoney Creek R.V. Park (data reviewed in April, 2008).

After combining the contaminant risk for bacteria and viruses with the natural susceptibility of the well, the overall vulnerability of the well to contamination is **Low**.

Nitrates and Nitrites

The contaminant risk for nitrates and nitrites is **Low** with septic systems, roads, and a campground contributing to the risk to the drinking water well.

The sampling history for the Stoney Creek R.V. Park well indicates that nitrates have been detected in the water within the last five years, with the highest concentration of 0.5 mg/l detected on 05/15/2007 (data reviewed in April, 2008).

After combining the contaminant risk for nitrates and nitrites with the natural susceptibility of the well, the overall vulnerability of the well to contamination is **Medium**.

Volatile Organic Chemicals

The contaminant risk for volatile organic chemicals is **Medium** with septic systems, heating oil tanks, roads, and a campground contributing to the risk to the drinking water well.

The drinking water at Stoney Creek R.V. Park has not been recently sampled for volatile organic chemicals (data reviewed in April, 2008).

After combining the contaminant risk for volatile organic chemicals with the natural susceptibility of the well, the overall vulnerability of the well to contamination is **Medium**.

Using the Source Water Assessment

This assessment of contaminant risks can be used as a foundation for local voluntary protection efforts as well as a basis for the continuous efforts on the part of Stoney Creek R.V. Park to protect public health. It is anticipated that Source Water Assessments will be updated every five years to reflect any changes in the vulnerability and/or susceptibility of Stoney Creek R.V. Park drinking water source.

REFERENCES


Alaska Department of Commerce, Community and Economic Development (ADCCED), Accessed 2008 [WWW document]. URL: http://www.commerce.state.ak.us/dca/commdb/CF_COMDB.htm

Freeze, R.A. and Cherry, J.A., 1979. Groundwater. Prentice-Hall, Englewood Cliffs, NJ.

United States Environmental Protection Agency (EPA), Accessed 2008 [WWW document]. URL: http://www.epa.gov/safewater/contaminants/index.html.

APPENDIX A

Stoney Creek R.V. Park Drinking Water Protection Area Location Map (Map A)




0

7,000

3,500

14,000



Stoney Creek RV Park PWS 249049.001 Appendix A Map A

APPENDIX B

Contaminant Source Inventory and Risk Ranking for Stoney Creek R.V. Park (Tables 1-4)

Contaminant Source Inventory for Stoney Creek RV Park

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Map Number	Comments
Septic systems (serves one single-family home)	R02	R02	А	С	31 inferred
Tanks, heating oil, residential (above ground)	R08	R08	А	С	31 assumed
Highways and roads, paved (cement or asphalt)	X20	X20	А	С	1 road
Campgrounds and RV Parks	X35	X35-01	А	С	
Septic systems (serves one single-family home)	R02	R02	В	С	37 inferred
Tanks, heating oil, residential (above ground)	R08	R08	В	С	37 assumed
Highways and roads, paved (cement or asphalt)	X20	X20	В	С	1 road

Table 2

Contaminant Source Inventory and Risk Ranking for

PWSID 249049.001

Stoney Creek RV Park Sources of Bacteria and Viruses

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Septic systems (serves one single-family home)	R02	R02	А	Low	С	31 inferred
Highways and roads, paved (cement or asphalt)	X20	X20	А	Low	С	1 road
Campgrounds and RV Parks	X35	X35-01	А	Low	С	
Septic systems (serves one single-family home)	R02	R02	В	Low	С	37 inferred
Highways and roads, paved (cement or asphalt)	X20	X20	В	Low	С	1 road

Table 3

Contaminant Source Inventory and Risk Ranking for

PWSID 249049.001

Stoney Creek RV Park Sources of Nitrates/Nitrites

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Septic systems (serves one single-family home)	R02	R02	А	Low	С	31 inferred
Highways and roads, paved (cement or asphalt)	X20	X20	А	Low	С	1 road
Campgrounds and RV Parks	X35	X35-01	А	Low	С	
Septic systems (serves one single-family home)	R02	R02	В	Low	С	37 inferred
Highways and roads, paved (cement or asphalt)	X20	X20	В	Low	С	1 road

Table 4

Contaminant Source Inventory and Risk Ranking for

PWSID 249049.001


Stoney Creek RV Park Sources of Volatile Organic Chemicals


Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Septic systems (serves one single-family home)	R02	R02	А	Low	С	31 inferred
Tanks, heating oil, residential (above ground)	R08	R08	А	Medium	С	31 assumed
Highways and roads, paved (cement or asphalt)	X20	X20	А	Low	С	1 road
Campgrounds and RV Parks	X35	X35-01	А	Low	С	
Septic systems (serves one single-family home)	R02	R02	В	Low	С	37 inferred
Tanks, heating oil, residential (above ground)	R08	R08	В	Medium	С	37 assumed
Highways and roads, paved (cement or asphalt)	X20	X20	В	Low	С	1 road

APPENDIX C

Stoney Creek R.V. Park Drinking Water Protection Area and Potential and Existing Contaminant Sources (Map C)

Public Water Well System for PWS #249049.001 Stoney Creek RV Park Showing Potential and Existing Sources of Contamination





Stoney Creek RV Park PWS 249049.001 Appendix C Map C