

# Hydrogeologic Susceptibility and Vulnerability Assessment for River's Edge Recreation Park Drinking Water Well, Sutton, Alaska

DRINKING WATER PROTECTION PROGRAM REPORT 103

---

September 2001

# Hydrogeologic Susceptibility and Vulnerability Assessment for River's Edge Recreation Park Drinking Water Well, Sutton, Alaska

By Catherine Baxter, B.E.S.T. Resource

---

DRINKING WATER PROTECTION PROGRAM REPORT 103

ALASKA DEPARTMENT OF ENVIRONMENTAL CONSERVATION: 2001

---

## CONTENTS

	Page		Page
Executive Summary	1	Inventory of Potential and Existing	
Introduction	1	Contaminant Sources	3
Description of the Matanuska – Susitna Valley, Alaska	1	Ranking of Contaminant Risks	3
River’s Edge Recreation Park Public Water Source	2	Vulnerability of Rivers Edge Recreation Park	
Assessment/Protection Area for River’s Edge		Drinking Water Source	4
Recreation Park Drinking Water Source	3	Summary	5
		References Cited	6

## TABLES

TABLE	1. Natural Susceptibility - Susceptibility of the Wellhead and Aquifer to Contamination	4
	2. Contaminant Risks	4
	3. Overall Vulnerability of River’s Edge Recreation Park Public Drinking Water Source to Contamination	5

## ILLUSTRATIONS

FIGURE	1. Index map showing the location of well assessment
--------	--

## APPENDICES

APPENDIX	A. River’s Edge Recreation Park Drinking Water Protection Area (Map 1)
	B. Contaminant Source Inventory for River’s Edge Recreation Park (Table 1)
	Contaminant Source Inventory and Risk Ranking for River’s Edge Recreation Park – Bacteria and Viruses (Table 2)
	Contaminant Source Inventory and Risk Ranking for River’s Edge Recreation Park – Nitrates/Nitrites (Table 3)
	Contaminant Source Inventory and Risk Ranking for River’s Edge Recreation Park – Volatile organic chemicals (Table 4)
	C. River’s Edge Recreation Park Drinking Water Protection Area and Potential and Existing Contaminant Sources (Map 2)
	D. Vulnerability Analysis for Contaminant Source Inventory and Risk Ranking for River’s Edge Recreation Park Public Drinking Water Source (Chart 1 – Chart 8 and Table 1 – Table 3)

# Hydrogeologic Susceptibility and Vulnerability Assessment for River's Edge Recreation Park Public Drinking Water Source, Palmer, Alaska

By Catherine Baxter, B.E.S.T. Resource

## Drinking Water Protection Program Alaska Department of Environmental Conservation

### EXECUTIVE SUMMARY

The River's Edge Recreation Park is a Class B (transient/noncommunity) drinking water source consisting of one well. Identified potential and current sources of contaminants for River's Edge Recreation Park include: one pit toilet and a RV dump station. This existing and potential source of contamination is considered a source of bacteria and viruses, nitrates and/or nitrites, and volatile organic chemicals. Overall, River's Edge Recreation Park public water source received a vulnerability rating of **Medium** for bacteria and viruses, nitrates and/or nitrites, and volatile organic chemicals.




Figure 1. Index map showing the location of well assessment

### INTRODUCTION

The purpose of this environmental assessment is to provide public water system owners/operators, communities, and local governments with information they can use to preserve the quality of Alaska's public drinking water supplies. This assessment was

completed for the River's Edge Recreation Park source of public drinking water. This source consists of one well in the Sutton area (Figure 1). This assessment, known under the Alaska Drinking Water Protection Program as the *Source Water Assessment*, has combined a review of the natural hydrogeologic sensitivity with potential and existing contaminant risks to arrive at an overall vulnerability of the drinking water source to contamination. This assessment has been completed as a basis for local voluntary protection efforts and to assist agencies in their efforts to reduce risk to this public drinking water supply.

### DESCRIPTION OF THE MATAMUSKA-SUSITNA VALLEY-AREA, ALASKA

#### Location

The Matanuska-Susitna Valley is part of the lowland lying about 50 miles north of Anchorage in south-central Alaska. The well described in this report is part of the Matanuska River Watershed. This study area is roughly bounded on the north by the Talkeetna Mountains; on the west by Wasilla Creek; on the south by the Knik River; and on the east by the Chugach Mountains. The area covers approximately 150 square miles.

#### Climate

The climate of the Matanuska-Susitna Valley is the result of a combination of marine and continental influences. The climate is somewhat transitional in that it does not experience large daily and annual temperature fluctuations like those experienced in the interior of Alaska nor does it experience high amounts of precipitation typified by gulf coast regions. Mean annual precipitation is approximately 15 inches per year. On the average, the Valley receives a total snow accumulation of 58 inches per year. Precipitation generally increased inland toward the Talkeetna Mountains where annual precipitation may exceed 60 inches. Mean daily temperature ranges from 67° F during July to 5° F in January [*Western Regional Climate Center, 2000*].

## **Physiography and Groundwater Conditions**

The Matanuska-Susitna Valley is surrounded by rugged mountains that rise abruptly from the valley floor. The Chugach Mountains at the southern edge of the valley reach altitudes greater than 6300 feet. These mountains are composed primarily of metamorphosed sedimentary marine and volcanic rocks, and greenstone of Mesozoic age. Along the northern edge of the valley, peaks in the Talkeetna Mountains reach altitudes of 3000 to 5000 feet. The Talkeetna Mountains are composed mainly of igneous rocks, chiefly granitic intrusives (Mesozoic?) and subordinate lavas and tuffs; Cretaceous and Tertiary sedimentary rocks form the south flank of the mountains. Although the altitude of the valley floor ranges from sea level at Knik Arm to 1000 feet at the base of Wishbone Hill, the local relief is commonly not more than 100 to 200 feet.

The Matanuska and Knik River's drain the area. These rivers are braided glacial outwash streams having wide floodplains. Drainage is poor in many interstream tracts resulting in large areas of swampy ground with shallow lakes occupying depressions.

The Matanuska-Susitna Valley is floored with unconsolidated deposits, chiefly glacial drift, that represents several episodes of glacial advances and retreats. The drift includes till, outwash stream deposits, and estuarine and lake deposits. Physiographic features formed by these deposits in or adjacent to the study area include end moraine, lateral moraines, eskers, crevasse fillings, and other pitted features, river terraces, outwash floodplains and an extensive estuarine flat (Trainer, 1960).

The glacial till and bedrock form aquifers of minor importance. The chief hydrologic significance of the till is in confining the artesian aquifer. Generally, the till is poorly permeable, although locally thin layers of sand may yield small quantities of water. Till that is present at or near the land surface in much of the area makes the acquisition of shallow groundwater difficult. The bedrock is poorly permeable. It yields water only from fractures, whose location and frequency cannot be easily predicted.

The chief aquifers are composed of outwash sand and gravel laid down by melt-water streams or in lakes. The outwash deposits are of two chief forms. The first consists of sheetlike deposits that lie just beneath the ground surface. These deposits range in thickness from a few feet to more than 100 feet. They typically rest on till or bedrock. The water in these deposits is unconfined. The other outwash deposits are buried

beneath till. They are known to be as much as 50 to 60 feet thick, and probably are considerably thicker in some places. They commonly contain confined, or artesian, groundwater. Well logs and data from pumping tests suggest that outwash sand and gravel form a continuous or nearly continuous sheet in an area of more than 10 square miles north and west of Palmer (Jakola et al, 1991).

Recharge of the groundwater is chiefly from precipitation but it is likely that only a small proportion of the annual precipitation reaches the water body. During very dry seasons conspicuous declines in of water levels occur in many wells. Along the mountain fronts, groundwater seeps from fractures in bedrock into the sediments. At these higher elevations, rain and snowmelt also enter the sediments. Lastly, aquifers may be recharged by streams where surface water percolates into surrounding permeable sediments (losing reaches of streams). This is the case for the water-table aquifers in the terrace south of Palmer and in the Bodenburg Butte area, which receive underground flow from the Matanuska River. Groundwater flow in the confined aquifers is generally from the north and north-northwest. The direction of groundwater flow in the upper unconfined aquifer is more variable due to the influence from surficial topography as well as its close connection with surface water bodies (Trainer, 1960).

## **RIVER'S EDGE RECREATION PARK PUBLIC WATER SOURCE**

River's Edge Recreation Park public water source is a Class B (transient/noncommunity) water source, which is privately owned and operated. The source consists of one well near the base of the Talkeetna Mountains at an elevation of 550 feet above sea level. The well is located north of the Glennallen Highway approximately 1 mile east of Sutton near Granite Creek. According to the well log, River's Edge Recreation Park does not appear to be grouted, but is functioning properly. The well penetrates gravel and silty sand with one 10-foot layer of gray clay from 30 to 40 feet below land surface. The well does not appear to be screened. The well had a static water level of 30 feet below land surface at the time of drilling (8/5/86).

The water from River's Edge Recreation Park consists of a hydropneumatic pressure tank, and jet pump. This water source operates 150 days per year. The River's Edge Recreation Park drinking water source collectively serves approximately 25 residents and non-residents through three service connections.

## **ASSESSMENT AND PROTECTION AREA FOR RIVER'S EDGE RECREATION PARK DRINKING WATER SOURCE**

The Drinking Water Protection and Assessment Area that has been established for River's Edge Recreation Park is the area that is most sensitive to contamination. This area has served as a basis for assessing the risk of the drinking water source to contamination. This zone around the drinking water source is the most critical area for the preservation of the quality of the drinking water for this source. For simplicity, this area will be known as your Drinking Water Protection Area and will serve as the area of focus for voluntary protection efforts.

Conceptually, groundwater enters the aquifer systems along the front range of the Talkeetna Mountains and flows toward Cook Inlet. An analytical calculation was used to calculate the size and shape of the area that contributes water to the well. The input parameters describing the attributes of the aquifer in this calculation were adopted from the well log and the recent Sanitary Survey. This analytical calculation was used as a guide in establishing the protection area for River's Edge Recreation Park. Additional methods were further employed to take into account any uncertainties in groundwater flow and aquifer characteristics to arrive at a meaningful and conservative protection area with respect to public health (Please refer to the Guidance Manual for Class B Public Water Systems for additional information).

The Drinking Water Protection Areas established for wells by the Alaska Department of Environmental Conservation (ADEC) are separated into zones. These zones correspond to a time-of-travel. Time-of-travel is the time required for water to move in the saturated zone of the ground from a specific point to the well. The Drinking Water Protection Areas for River's Edge Recreation Park contains four zones, Zone A, Zone B, Zone C and Zone D (Map 1, Appendix A). Zone A corresponds to the area between the well and the distance equal to  $\frac{1}{4}$  of the distance of the 2-year time-of-travel. Depending on where a contaminant source is located within Zone A, travel time for a contaminant to the well may be on the order of several days to several hours. Zone A also extends down gradient from the well to take into account the area of the aquifer that is influenced by pumping of the well.

The Zone B protection area for River's Edge Recreation Park corresponds to a time-of-travel of less than two years and extends toward base of the Talkeetna Mountains. Zone C protection area corresponds to a

time-of-travel of greater than 2 years and less than 5 years. Zone D corresponds to a time-of-travel of greater than 5 years and less than 10 years.

## **INVENTORY OF POTENTIAL AND EXISTING CONTAMINANT SOURCES**

The Drinking Water Protection Program has completed an inventory of potential and existing sources of contamination within River's Edge Recreation Park Drinking Water Protection Area. This survey was completed through a search of agency records and other publicly available information.

Potential sources of contamination to drinking water supplies cover a wide range of categories and types. Potential drinking water contaminants are found within agricultural, residential, commercial, and industrial areas, but can also occur within areas that have little or no development.

For the basis of this assessment and all Class B public water system assessments, three categories of drinking water contaminants were inventoried. They include:

- Bacteria and viruses
- Nitrates and/or nitrites
- Volatile organic chemicals.

Table 1 in Appendix C lists the Contaminant Source Inventory for River's Edge Recreation Park. Inventoried potential sources of contamination within Zone A were attributed to waste disposal. Zones B, C and D contain only natural wilderness and were not considered in determining the vulnerability of this drinking water source to contamination. Below is a summary of the contaminant sources inventoried within the River's Edge Recreation Park protection area:

- Pit Toilet
- RV Dump Station.

This potential contaminant source presents risks for all three categories of drinking water contaminants for River's Edge Recreation Park drinking water source.

## **RANKING OF CONTAMINANT RISKS**

Potential and existing sources of contamination have been identified, sorted, and ranked according to what type and level of risk they represent. Ranking of contaminant risks for a "potential" or "existing" source of contamination is a function of toxicity and volumes of specific contaminants associated with that source. Contaminant risks are further a function of the number

and density of those types of contaminant sources as well as the proximity of those sources to the well (Appendices B & C).

**VULNERABILITY OF RIVER’S EDGE RECREATION PARK DRINKING WATER SOURCES**

Vulnerability of a drinking water source to contamination is a combination of two factors:

- Natural susceptibility; and
- Contaminant risks.

Each of the three categories of drinking water contaminants has been analyzed and an overall vulnerability score of 0 to 100 is ultimately assigned:

$$\begin{aligned}
 &\text{Natural Susceptibility (0 – 50 points)} \\
 &\quad + \\
 &\quad \text{Contaminant Risks (0 – 50 points)} \\
 &\quad = \\
 &\quad \text{Vulnerability of the} \\
 &\quad \text{Drinking Water Source to Contamination (0 – 100).}
 \end{aligned}$$

A score for the Natural Susceptibility is achieved by analyzing the properties of the well and the aquifer.

$$\begin{aligned}
 &\text{Susceptibility of the Wellhead (0 – 25 Points)} \\
 &\quad + \\
 &\quad \text{Susceptibility of the Aquifer (0 – 25 Points)} \\
 &= \text{Natural Susceptibility (Susceptibility of the Well)} \\
 &\quad \text{(0 – 50 Points)}
 \end{aligned}$$

River’s Edge Recreation Park is completed in a confined or semi-confined aquifer setting. The well penetrates 51 feet of sand and gravel with one clay-bearing layer. This 10-foot clay layer, encountered at 30 feet below land surface, may provide a protective barrier for the movement of contaminants in the subsurface. However, near the base of the Talkeetna Mountains, the clay and till layers tend to be discontinuous and thin toward the mountains. Therefore, contaminants that enter the subsurface near the base of the mountains may enter the confined aquifer uninhibited by any protective layer. This well does not appear to be properly grouted as indicated previously from information obtained from ADEC records. The absence of grouting can promote the transport of contaminants along the well casing.

Combining the susceptibilities of the wellhead and the aquifer to contamination leads to a score (0 – 50 points) and rating of overall Susceptibility (Appendix D). Table 1 shows the overall Susceptibility score and rating for River’s Edge Recreation Park.

**Table 1. Natural Susceptibility - Susceptibility of the Wellhead and Aquifer to Contamination**

	Score	Rating
Susceptibility of the Wellhead	10	Medium
Susceptibility of the Aquifer	23	Very High
Natural Susceptibility	33	High

Contaminant risks to a drinking water source depend on the type, number or density, and distribution of contaminant sources. One Pit Toilet and one RV Dump Station contribute the highest and only identified risk for potential contamination to the River’s Edge Recreation Park source of public drinking water.

A score (0 – 50 points) and rating of Contaminant Risks (Appendix D) is assigned based on the findings of the Contaminant Source Inventory (Appendix B - Table 1 – Table 7). This portion of the analysis examines any existing or historical contamination that has been detected at the drinking water source through routine sampling. It also reviews contamination that has or may have occurred but has not arrived or been detected at the well. Table 2 summarizes the Contaminant Risks for each category of drinking water contaminants.

**Table 2. Contaminant Risks**

Contaminant Risks	Score	Rating
Bacteria and Viruses	12	Low
Nitrates and/or Nitrites	16	Low
Volatile Organic Chemicals	10	Low

Appendix D contains eight charts, which together form the ‘Vulnerability Analysis’ for a source water assessment for a public drinking water source. Chart 1 analyzes the ‘Susceptibility of the Wellhead’ to contamination by looking at the construction of the well and its surrounding area. Chart 2 analyzes the ‘Susceptibility of the Aquifer’ to contamination by looking at the naturally occurring attributes of the water source and influences on the groundwater system that might lead to contamination. Chart 3 analyzes ‘Contaminant Risks’ for the drinking water source with

respect to bacteria and viruses. The ‘Contaminant Risks’ portion of the analysis considers potential sources of contaminants as well as a review of contamination that has or may have occurred but has not arrived or been detected at the well. Lastly, Chart 4 contains the ‘Vulnerability Analysis for Bacteria and Viruses’. Charts 5 through 8 contain the Contaminant Risks and Vulnerability Analysis for nitrates and nitrites and volatile organic chemicals, respectively.

Vulnerability of the drinking water source to contamination is the combination of susceptibility of the aquifer and the well with contaminant risks. Table 3 contains the overall vulnerability scores (0 – 100) and ratings for each of the three categories of drinking water contaminants (Appendix D). Note: scores are rounded off to the nearest five.

**Table 3. Overall Vulnerability of River’s Edge Recreation Park Public Drinking Water Source to Contamination by Category**

Category	Score	Rating
Bacteria and Viruses	45	Medium
Nitrates and Nitrites	50	Medium
Volatile Organic Chemicals	40	Medium

Tables 2 through 4 in Appendix B contain the ranking of potential and existing sources of contamination with respect to bacteria and viruses, nitrates and/or nitrites, and volatile organic chemicals.

The Pit Toilet and RV Dump Station in Zone A are the only factors in determining contaminant risks for all categories of contaminants (See “Overall Rank after Analysis” in Table 2 – 4 of Appendix B).

Bacteria and Viruses were not detected in the source waters of River’s Edge Recreation Park.

Sampling history of River’s Edge Recreation Park source waters indicate low concentrations of nitrate (See Chart 6 – Contaminant Risks for Nitrates/Nitrites in Appendix D). Existing nitrate contamination is approximately 1% of the allowable limit (MCL) for this contaminant. Due to the high solubility and weak retention by soil, nitrates are very mobile in soil, moving at approximately the same rate as water. The current nitrate concentration in River’s Edge Recreation Park remains at safe levels with respect to human health.

Overall, contaminant risks for the nitrate/nitrite category are medium due to the pit toilet and RV dump station present up gradient from the well. Combining potential nitrate and/or nitrite contamination risk with the susceptibility of the well yields an overall medium vulnerability to contamination in this category.

Volatile Organic Chemicals were not detected in the source waters of River’s Edge Recreation Park.

## SUMMARY

A *Source Water Assessment* has been completed for the River’s Edge Recreation Park source of public drinking water. The overall vulnerability of this source to contamination is medium for bacteria and viruses, high for nitrates and/or nitrites, and medium for volatile organic chemicals. This assessment of contaminant risks can be used as a foundation for local voluntary protection efforts as well as a basis for the continuous efforts on the part of the River’s Edge Recreation Park to protect public health. It is anticipated that *Source Water Assessments* will be updated every five years to reflect any changes in the vulnerability and/or susceptibility of the public drinking water source.



## REFERENCES CITED

Jakola, J.B., Munter, J.A., and Evans, J.G., 1991, Ground-water resources of the Palmer-big Lake area, Alaska: a conceptual model. Division of Geological & Geophysical Surveys Reported of Investigations 90-4, State of Alaska Department of Natural Resources, Fairbanks, AK.


Trainer, F.W., 1960, Geology and Groundwater Resources, Matanuska Valley, Alaska, U.S. Geological Survey Water Supply Paper 1494 U.S. Printing Office, Washington, D.C.










Western Regional Climate Center, 2000, August 24, Web extension to the *Western Regional Climate Center*  
[WWW document]. URL [http://www.uaa.alaska.edu/enri/ascc\\_web/ascc\\_home.html](http://www.uaa.alaska.edu/enri/ascc_web/ascc_home.html) .

## **APPENDIX A**

### **River's Edge Recreation Park Drinking Water Protection Area**

# River's Edge Recreation Park (PWSID 227377) Drinking Water Protection Areas



-  North
-  Public Water System Wells
-  Lakes and Rivers
-  Zone A Protection Area  
Several Months to 2 Years  
Travel Time
-  Zone B Protection Area  
Less Than 2 Years  
Travel Time
-  Zone C Protection Area  
2 to 5 Years Travel Time
-  Zone D Protection Area  
5 to 10 Years Travel Time
-  Roads
-  Mat-Su Borough Parcel  
Boundaries



## Map 1

## **APPENDIX B**

### **Contaminant Source Inventory and Risk Ranking for River's Edge Recreation Park**

Table 1

**Contaminant Source Inventory for  
River's Edge Recreation Park**

PWSID 227377.001

<b>Contaminate Source Category</b>	<b>Contaminant Source ID</b>	<b>CS ID Tag</b>	<b>Zone</b>	<b>Location</b>	<b>Map</b>	<b>Comments</b>
Pit Toilet	D16	D16-1	A	Located approx. 225' north of well	2	
RV Dump Station	D18	D18-1	A	Located approx. 325' northeast of well	2	

Table 2

**Potential and Existing Sources of Contamination for  
River's Edge Recreartion Park  
Bacterias and Viruses**

PWSID 227377.001

<b>Contaminant Source Category</b>	<b>Contaminant Source ID</b>	<b>CS ID Tag</b>	<b>Zone</b>	<b>Risk Ranking for Analysis</b>	<b>Overall Rank for Analysis</b>	<b>Location</b>	<b>Map</b>	<b>Comments</b>
Pit Toilet	D16	D16-1	A	Medium	1	Located approx. 225' north of well	2	
RV Dump Station	D18	D18-1	A	Low	2	Located approx. 325' northeast of well	2	

Table 3

**Potential and Existing Sources of Contamination for  
River's Edge Recreation Park  
Nitrates and Nitrites**

PWSID 227377.001

<b>Contaminant Source Category</b>	<b>Contaminant Source ID</b>	<b>CS ID Tag</b>	<b>Zone</b>	<b>Risk Ranking for Analysis</b>	<b>Overall Rank for Analysis</b>	<b>Location</b>	<b>Map</b>	<b>Comments</b>
Pit Toilet	D16	D16-1	A	Medium	1	Located approx. 225' north of well	2	
RV Dump Station	D18	D18-1	A	Low	2	Located approx. 325' northeast of well	2	

Table 4

**Potential and Existing Sources of Contamination for  
River's Edge Recreation Park  
Volatile Organic Chemicals (VOCs)**

PWSID 227377.001



<b>Contaminant Source Category</b>	<b>Contaminant Source ID</b>	<b>CS ID Tag</b>	<b>Zone</b>	<b>Risk Ranking for Analysis</b>	<b>Overall Rank for Analysis</b>	<b>Location</b>	<b>Map</b>	<b>Comments</b>
Pit Toilet	D16	D16-1	A	Low	1	Located approx. 225' north of well	2	
RV Dump Station	D18	D18-1	A	Low	2	Located approx. 325' northeast of well	2	



## **APPENDIX C**

### **River's Edge Recreation Park Drinking Water Protection Area and Potential & Existing Contaminant Sources**

# Drinking Water Protection Areas Potential & Existing Sources of Contamination for River's Edge Recreation Park



- Public Water System Wells
- Zone A Protection Area  
Several Months to 2 Years  
Travel Time
- Zone B Protection Area  
Less Than 2 Years  
Travel Time
- ▬ Roads
- ▭ Mat-Su Borough Parcel  
Boundaries
- Pit toilet (D16)
- RV Dump Station (D18)



PWSID 227377.001




## Map 2

Prepared by: B.E.S.T. Resource

## **APPENDIX D**


### **Vulnerability Analysis for River's Edge Recreation Park Public Drinking Water Source**

**Chart 1. Susceptibility of the wellhead – River’s Edge Recreation Park**





<u>Wellhead Susceptibility Ratings</u>	
20 to 25 pts	very high
15 to < 20 pts	high
10 to < 15 pts	medium
< 10	low

**Chart 2. Susceptibility of the aquifer – River's Edge Recreation Park**



Recharge (16-20 inches per year, base of Talkeetna Mountains, with silty, gravelly sand) 4/10 = 4 Points  
 Depth to static water level (30 feet) 4/10 = 4 Points  
 Protectiveness of the Vadose Zone Total = 8 Points



Unconfined 15/15 = 10 Points  
 Density of boreholes/wells 15/15 = 5 Point  
 Degree of Confinement Total = 15 Points


**15 Points**

**Very High**

Aquifer Susceptibility Ratings	
20 to 25 pts	very high
15 to < 20 pts	high
10 to < 15 pts	medium
< 10	low

- Protectiveness of the Vadose Zone**
  - net recharge (function of precipitation, slope of land surface, & permeability of soils) [0-10 pts: 50%]
  - depth to water table (unconfined aquifer) or top of confining layer (confined aquifer) [interpolate linearly: 100' - 20', 0-5 pts; 20' - 0', 5-10 pts; 50% weight]
- Degree of Confinement**
  - confined versus unconfined aquifer [confined:  $K \leq 10^{-6}$  cm/s, minimum thickness of at least one layer = 20 ft, interpolate linearly 100' - 20', 0 - 10 pts; unconfined = 15 pts; 65%]
  - density of boreholes and wells penetrating the confining layer (confined aquifer) or the water table (unconfined aquifer) [confined: 0 - 15 pts; unconfined = 15 pts; 35% weight]

Chart 3. Contaminant risks for River's Edge Recreation Park – Bacteria & Viruses



1 Pit toilet = 10+ Lows  
 1 RV Dump Station = 1 Low  
 Total = 10+ 1 Lows

+10 Points

\* Exclude sources of existing contamination, if known

Chart 3. Contaminant risks for River's Edge Recreation Park – Bacteria & Viruses (Continued)







Chart 3. Contaminant risks for River's Edge Recreation Park – Bacteria & Viruses (Continued)





**Table 1. Risk Matrix for Contaminant Sources for River’s Edge Recreation Park – Bacteria & Viruses**

**Level of Risk Associated with the Highest Risk Sources**

<b>Next Highest Risk Sources(s)</b>	1 Pit toilet, 1 RV Dump Station	<b>LOW</b> 10 pts	<b>MEDIUM</b> 20 pts	<b>HIGH</b> 30 pts	<b>VERY HIGH</b> 40 pts
	<b>Low</b>	≥ 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	—
	<b>Medium</b>	—	≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
	<b>High</b>	—	—	1 source + 10 pts	≥ 2 sources + 10 pts
	<b>Very High</b>	—	—	—	1 source + 10 pts

**Chart 4. Vulnerability analysis for River's Edge Recreation Park– Bacteria & Viruses**






Chart 5. Contaminant risks for River's Edge Recreation Park– Nitrates and Nitrites




Chart 5. Contaminant risks for River's Edge Recreation Park– Nitrates and Nitrites (Continued)






Chart 5. Contaminant risks for River's Edge Recreation Park– Nitrates and Nitrites (Continued)




**Table 2. Risk Matrix for Contaminant Sources for River’s Edge Recreation Park– Nitrates and Nitrites**

**Level of Risk Associated with the Highest Risk Sources**

1 PIT TOILET, 1 RV DUMP STATION	<b>LOW</b> 10 pts	<b>MEDIUM</b> 20 pts	<b>HIGH</b> 30 pts	<b>VERY HIGH</b> 40 pts
<b>Low</b>	> 10 sources + 10 pts	> 10 sources + 5 pts	> 20 sources + 5 pts	---
<b>Medium</b>	---	> 2 sources + 5 pts	> 5 sources + 5 pts	> 10 sources + 5 pts
<b>High</b>	---	---	1 source + 10 pts	> 2 sources + 10 pts
<b>Very High</b>	---	---	---	1 source + 10 pts

**Chart 6. Vulnerability analysis for River's Edge Recreation Park– Nitrates and Nitrites**



**Chart 7. Contaminant risks for River's Edge Recreation Park – Volatile Organic Chemicals**






Chart 7. Contaminant risks for River's Edge Recreation Park – Volatile Organic Chemicals (Continued)





Chart 7. Contaminant risks for River's Edge Recreation Park – Volatile Organic Chemicals (Continued)



**Table 3. Risk Matrix for Contaminant Sources for River’s Edge Recreation Park – Volatile Organic Chemicals**

**Level of Risk Associated with the Highest Risk Sources**

<b>Next Highest Risk Sources(s)</b>	No known existing or potential contaminate sources	<b>LOW</b> 10 pts	<b>MEDIUM</b> 20 pts	<b>HIGH</b> 30 pts	<b>VERY HIGH</b> 40 pts
	<b>Low</b>	> 10 sources + 10 pts	> 10 sources + 5 pts	> 20 sources + 5 pts	---
	<b>Medium</b>	---	> 2 sources + 5 pts	> 5 sources + 5 pts	> 10 sources + 5 pts
	<b>High</b>	---	---	1 source + 10 pts	> 2 sources + 10 pts
	<b>Very High</b>	---	---	---	1 source + 10 pts

**Chart 8. Vulnerability analysis for River's Edge Recreation Park– Volatile Organic Chemicals**

