



Source Water Assessment

A Hydrogeologic Susceptibility and Vulnerability Assessment for Leonard's Landing & Lodge, Yakutat, Alaska PWSID #130705

DRINKING WATER PROTECTION PROGRAM REPORT NO. 728

Alaska Department of Environmental Conservation

Source Water Assessment for Leonard's Landing & Lodge Yakutat, Alaska PWSID #130705

DRINKING WATER PROTECTION PROGRAM REPORT NO. 728

The Drinking Water Protection Program (DWPP) is producing Source Water Assessments in compliance with the Safe Drinking Water Act Amendments of 1996. Each assessment includes a delineation of the source water area, an inventory of potential and existing contaminant sources that may impact the water, a risk ranking for each of these contaminants, and an evaluation of the potential vulnerability of these drinking water sources.

These assessments are intended to provide public water systems owners/operators, communities, and local governments with the best available information that may be used to protect the quality of their drinking water. The assessments combine information obtained from various sources, including the U.S. Environmental Protection Agency, Alaska Department of Environmental Conservation (ADEC), public water system owners/operators, and other public information sources. The results of this assessment are subject to change if additional data becomes available. It is anticipated this assessment will be updated every five years to reflect any changes in the vulnerability and/or susceptibility of public drinking water source. If you have any additional information that may affect the results of this assessment, please contact the Program Coordinator of DWPP, (907) 269-7521.

CONTENTS

SECTION	Executive Summary Leonard's Landing & Lodge Public Drinking Water System Leonard's Landing & Lodge Drinking Water Protection Area Inventory of Potential and Existing Contaminant Sources Ranking of Contaminant Risks Vulnerability of Leonard's Landing & Lodge Drinking Water System References	Page 1 1 1 2 2 2 5
	TABLES	
TABLE	 Definition of Zones Susceptibility Contaminant Risks Overall Vulnerability 	2 3 3 3
	APPENDICES	
APPENDIX	 A. Leonard's Landing & Lodge Drinking Water Protection Area (Map 1) B. Contaminant Source Inventory for Leonard's Landing & Lodge (Table 1) Contaminant Source Inventory and Risk Ranking for Leonard's Landing & and Viruses (Table 2) Contaminant Source Inventory and Risk Ranking for Leonard's Landing & Nitrates/Nitrites (Table 3) Contaminant Source Inventory and Risk Ranking for Leonard's Landing & Volatile Organic Chemicals (Table 4) 	Lodge –
	C. Leonard's Landing & Lodge Drinking Water Protection Area and Potential and Existing Sources of Contamination (Map 2)	
	D. Vulnerability Analysis for Contaminant Source Inventory and Risk Ranking Leonard's Landing & Lodge Public Drinking Water Source (Charts 1 –	

Drinking Water Protection Program Alaska Department of Environmental Conservation

EXECUTIVE SUMMARY

The public water system for Leonard's Landing & Lodge is a Class B (transient/non-community) water system consisting of two wells. Leonard's Landing & Lodge is located northwest of Yakutat, Alaska. The wellhead received a susceptibility rating of Low and the aquifer received a susceptibility rating of Very High. Combining these two ratings produces a Medium rating for the natural susceptibility of the well. Identified potential and current sources of contaminants for Leonard's Landing & Lodge public drinking water source include: dirt/gravel highways and roads, and a septic system. This identified potential and existing source of contamination is considered a source of bacteria and viruses, nitrates and/or nitrites, and volatile organic chemicals. Overall, the public water source for Leonard's Landing & Lodge received a vulnerability rating of Low for both bacteria and viruses; Medium for nitrates and nitrites, and Low for volatile organic chemicals.

LEONARD'S LANDING & LODGE PUBLIC DRINKING WATER SYSTEM

Leonard's Landing & Lodge public water system is a Class B (transient/non-community) water system. The system consists of two wells that are currently in service at the Lodge northwest of Yakutat, Alaska (See Map 1 of Appendix A). Yakutat is located on the Gulf of Alaska coast, where southeastern Alaska joins the major body to Alaska to the west. Yakutat is 225 miles northwest of Juneau. The population of Yakutat is approximately 800.

Yakutat averages about 150 inches of precipitation per year; and approximately 200 inches of snow. The groundwater aquifers underlying the area are recharged through the infiltration of precipitation and surface water. Groundwater aquifers in the region generally occur in the fractured bedrock and unconsolidated sediments deposited by glaciers and/or rivers. The elevation for Yakutat is near sea level. The Leonard's Landing & Lodge area topography is near sea level.

According to a Sanitary Survey dated June 9, 1999, Well No. 1 was installed in 1995, and Well No. 2 was installed pre-1989. Both wells were installed at a depth

of 40 feet with 8 inch casing. It is assumed that the length of the well screens are 10 feet.

The Survey for the water system indicates the land surface is sloped away from the well, which would provide adequate surface water drainage. The wells are grouted according to ADEC regulations. Proper grouting provides added protection against contaminants traveling along the well casing and into source waters.

Well No. 1 operates seasonally, from March through October, and Well No. 2 operates year round. Each well supplies one half of the system, serving approximately 1 resident and 30 non-residents through two service connections.

LEONARD'S LANDING & LODGE DRINKING WATER PROTECTION AREA

In order to evaluate whether a drinking water source is at risk, we must first evaluate what are the most likely pathways for surface contamination to reach the groundwater. These areas are determined by looking at the characteristics of the soil, groundwater, aquifer, and well.

The most probable area for contamination to reach the drinking water well is the area that contributes water to the well, the groundwater recharge area. This area is designated as the Drinking Water Protection Area (DWPA). Because releases of contaminants within the DWPA are most likely to impact the drinking water well, this area will serve as the focus for voluntary protection efforts.

An analytical calculation was used to determine the size and shape of the DWPA. The input parameters describing the attribute of the aquifer in this calculation were adopted from the U.S. Geological Survey (Patrick, Brabets, and Glass, 1989), and State of Alaska Department of Water Resources (Jokela, et. al., 1991). Additional methods were also used to take into account any uncertainties in groundwater flow and aquifer characteristics to arrive at a meaningful DPWA (Please refer to the Guidance Manual for Class B Public Water Systems for additional information).

The DWPAs established for wells by the ADEC are usually separated into four zones. These zones correspond to differences in the time-of-travel (TOT) of the water moving through the aquifer to the well.

The TOT for contaminants within the water varies and is dependent on the physical and chemical characteristics of each contaminant. The following is a summary of the four protection area zones for wells and the calculated time-of-travel for each:

Table 1. Definition of Zones

Zone	Definition
A	¹ / ₄ the distance for the 2-yr. time-of-travel
В	Less than the 2 year time-of-travel
C	Less Than the 5 year time-of-travel
D	Less than the 10 year time-of-travel

The DWPA for Leonard's Landing & Lodge extends over a mile to the northwest of the well. Development in the vicinity of the well is limited to only Zone A (See Map 1 of Appendix A).

INVENTORY OF POTENTIAL AND EXISTING CONTAMINANT SOURCES

The Drinking Water Protection Program has completed an inventory of potential and existing sources of contamination within the Leonard's Landing & Lodge DWPA. This inventory was completed through a search of agency records and other publicly-available information. Potential sources of contamination to the drinking water aquifer include a wide range of categories and types. Potential drinking water contaminants are found within agricultural, residential, commercial, and industrial areas, but can also occur within areas that have little or no development.

For the basis of all Class B public water system assessments, three categories of drinking water contaminants were inventoried. They include:

- Bacteria and viruses;
- Nitrates and/or nitrites;
- Volatile organic chemicals

The sources summarized in Table 1 of Appendix B.

RANKING OF CONTAMINANT RISKS

Once the potential and existing sources of contamination have been identified, they are assigned a ranking according to what type and level of risk they represent. Ranking of contaminant risks for a "potential" or "existing" source of contamination is a function of toxicity and volumes of specific

contaminants associated with that source. Rankings include:

- Low:
- Medium;
- High; and
- Very High.

The TOT for contaminants within the water varies and is dependent on the physical and chemical characteristics of each contaminant. Bacteria and Viruses are only inventoried in Zones A and B because of their short life span. Only "Very High" and "High" rankings are inventoried within the outer Zone D due to the probability of contaminant dilution by the time the contaminants get to the well.

Tables 2 through 4 in Appendix B contain the ranking of potential and existing sources of contamination with respect to bacteria and viruses, nitrates and/or nitrites, and volatile organic chemicals.

VULNERABILITY OF LEONARD'S LANDING & LODGE DRINKING WATER SYSTEM

Vulnerability of a drinking water source to contamination is a combination of two factors:

- Natural susceptibility; and
- Contaminant risks.

Appendix C contains eight charts, which together form the 'Vulnerability Analysis' for a source water assessment for a public drinking water source. Chart 1 analyzes the 'Susceptibility of the Wellhead' to contamination by looking at the construction of the well and its surrounding area. Chart 2 analyzes the 'Susceptibility of the Aquifer' to contamination by looking at the naturally occurring attributes of the water source and influences on the groundwater system that might lead to contamination. Chart 3 analyzes 'Contaminant Risks' for the drinking water source with respect to bacteria and viruses. The 'Contaminant Risks' portion of the analysis considers potential sources of contaminants as well as a review of contamination that has or may have occurred, but has not arrived or been detected at the well. Lastly, Chart 4 contains the 'Vulnerability Analysis for Bacteria and Viruses'. Charts 5 through 8 contain the Contaminant Risks and Vulnerability Analyses for nitrates and nitrites and volatile organic chemicals, respectively.

A score for the Natural Susceptibility is reached by considering the properties of the well and the aquifer.

Susceptibility of the Wellhead (0 – 25 Points) (Chart 1 of Appendix C)

+

Susceptibility of the Aquifer (0 – 25 Points) (Chart 2 of Appendix C)

=

Natural Susceptibility (Susceptibility of the Well) (0-50 Points)

A ranking is assigned for the Natural Susceptibility according to the point score:

Natural Susceptibility Ratings

40 to 50 pts	Very High
30 to < 40 pts	High
20 to < 30 pts	Medium
< 20 pts	Low

We assume the well for the Leonard's Landing & Lodge is completed in an unconfined aquifer. Because unconfined aquifers are recharged by surface water and precipitation that migrates downward from the surface, contaminants at the surface have the potential to adversely impact this aquifer. Table 2 shows the Susceptibility scores and ratings for Leonard's Landing & Lodge

Table 2. Susceptibility

	Score	Rating
Susceptibility of the		
Wellhead	0	Low
Susceptibility of the		
Aquifer	25	Very High
Natural Susceptibility	25	Medium

Contaminant risks to a drinking water source depend on the type, number or density, and distribution of contaminant sources. This score has been derived from an examination of existing and historical contamination that has been detected at the drinking water source through routine sampling. It also evaluates potential sources of contamination. Flow charts are used to assign a point score, and ratings are assigned in the same way as for the natural susceptibility:

Contaminant Risk Ratings

40 to 50 pts	Very High
30 to < 40 pts	High
20 to < 30 pts	Medium
< 20 pts	Low

Table 3 summarizes the Contaminant Risks for each category of drinking water contaminants.

Table 3. Contaminant Risks

Category	Score	Rating
Bacteria and Viruses	4	Low
Nitrates and/or Nitrites	15	Low
Volatile Organic Chemicals	12	Low

Finally, an overall vulnerability score is assigned for each water system by combining each of the contaminant risk scores with the natural susceptibility score:

Natural Susceptibility (0 - 50 points)

+

Contaminant Risks (0 - 50 points)

=

Vulnerability of the Drinking Water Source to Contamination (0 - 100).

Again, rankings are assigned according to a point score:

Overall Vulnerability Ratings

80 to 100 pts	Very High
60 to < 80 pts	High
40 to < 60 pts	Medium
< 40 pts	Low

Table 4 contains the overall vulnerability scores (0 - 100) and ratings for each of the three categories of drinking water contaminants. Note: scores are rounded off to the nearest five.

Table 4. Overall Vulnerability

Category	Score	Rating
Bacteria and Viruses	30	Low
Nitrates and Nitrites	40	Medium
Volatile Organic Chemicals	35	Low

Bacteria and Viruses

The contaminant risk for bacteria and viruses is **Low** with the dirt/gravel highways and roads located within Zone A representing the risk to the drinking water well (See Chart 3 – Contaminant Risks for Bacteria and Viruses in Appendix C).

Only a small amount of bacteria and viruses are required to endanger public health. Sampling results indicates there were no consecutive positive tests for Bacteria and viruses. However, after combining the contaminant risks with the overall natural susceptibility of the well, the vulnerability of the well to contamination by bacteria and viruses is **Low**.

Nitrates and Nitrites

The contaminant risk for nitrates and nitrites is **Low** with the dirt/gravel highways and roads representing the risk to this source of public drinking water (See Chart 5 - Contaminant Risks for Nitrates and/or Nitrites in Appendix C).

Sampling history for Leonard's Landing & Lodge well indicates that nitrates have been detected in the water, but only in very low concentrations (most recently at 0.16 mg/L on 4/30/2002) or 2% of the Maximum Contaminant Level (MCL). The MCL is the maximum level of contaminant that is allowed to exist in drinking water and still be consumed by humans without harmful health effects. Due to the high solubility and weak retention by soil, nitrates are very mobile, moving at approximately the same rate as water.

After combining the contaminant risk for nitrates and nitrites with the natural susceptibility of the well, the overall vulnerability of the well to contamination by nitrates and nitrites is **Medium**.

Volatile Organic Chemicals

The contaminant risk for volatile organic chemicals is **Low** with the dirt/gravel highways and roads creating the only known risk for volatile organic chemicals (See Chart 7 – Contaminant Risks for Volatile Organic Chemicals in Appendix C).


There are no recent sample data available for the drinking water at Leonard's Landing & Lodge for volatile organic chemicals. However, after combining the contaminant risk for volatile organic chemicals with the natural susceptibility of the well, the overall vulnerability of the well to contamination by volatile organic chemicals is **Low**.

REFERENCES

- Alaska Department of Community and Economic Development (ADCED), 2002 [WWW document]. URL http://www.dced.state.ak.us/mra/CF_BLOCK.cfm.
- Alaska Geospatial Data Clearinghouse, 2003. URL: http://agdc.usgs.gov/data/datasets.html.
- Gehrels, G.E., Berg, H.C., Geologic Map of Southeastern Alaska: U.S. Geological Survey Map (scale 1:600,000), Map I-1867, 1sheet.
- Jokela, J.B., Munter, J.A., and Evans, J.G., 1991, Ground-water resources of the Palmer-Big Lake area, Alaska: a conceptual model. Division of Geological &Geophysical Surveys Reports of Investigations 90-4, State of Alaska Department of Natural Resources, Fairbanks, AK.
- King, P.B., compiler, 1969, Tectonic map of North America: US Geological Survey Map, (scale 1:5,000,000) 2 sheets.
- Patrick, L.D., Brabets, T.P., and Glass, R.L., 1989, Simulation of ground-water flow at Anchorage, Alaska: US Geological Survey Water-Resources Investigations Report 88-4139, 41p.
- United States Environmental Protection Agency (EPA), 2002 [WWW document]. URL: http://www.epa.gov/safewater/mcl.html.

APPENDIX A

Leonard's Landing & Lodge
Drinking Water Protection Area Location Map
(Map 1)



APPENDIX B

Contaminant Source Inventory and Risk Ranking for Leonard's Landing & Lodge (Tables 1-4)

Table 1

Contaminant Source Inventory for Leonard's Landing & Lodge

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Map Number	Comments
Highways and roads, dirt/gravel	X24	X24-1	A	2	Leonard's Landing & Lodge
Septic systems (serves one single-family home)	R02	R02-1-16	D	2	Sixteen single-family septic systems in Zone D
Water supply wells	W09	W09-1-16	D	2	Sixteen private water supply wells in Zone D
Highways and roads, dirt/gravel	X24	X24-2-5	D	2	Four roads throughout Zone D

Contaminant Source Inventory and Risk Ranking for

Table 2

PWSID 130705.001

Leonard's Landing & Lodge Sources of Bacteria and Viruses

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Highways and roads, dirt/gravel	X24	X24-1	A	Low	2	Leonard's Landing & Lodge

PWSID 130705.001

Contaminant Source Inventory and Risk Ranking for

Leonard's Landing & Lodge Sources of Nitrates/Nitrites

Table 3

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Highways and roads, dirt/gravel	X24	X24-1	A	Low	2	Leonard's Landing & Lodge

Contaminant Source Inventory and Risk Ranking for

Table 4


PWSID 130705.001

Leonard's Landing & Lodge Sources of Volatile Organic Chemicals

Contaminant Source Type	Contaminant Source ID	CS ID tag	Zone	Risk Ranking for Analysis	Map Number	Comments
Highways and roads, dirt/gravel	X24	X24-1	A	Low	2	Leonard's Landing & Lodge

APPENDIX C

Leonard's Landing & Lodge
Drinking Water Protection Area Location Map and Potential and
Existing Sources of Contamination
(Map 2)



APPENDIX D

Vulnerability Analysis for Leonard's Landing & Lodge Public Drinking Water Source (Charts 1-8)

Chart 1. Susceptibility of the wellhead - Leonard's Landing





Chart 2. Susceptibility of the aquifer - Leonard's Landing







Chart 3. Contaminant risks for Leonard's Landing - Bacteria & Viruses NO Initial assessment of risk posed by Are there sufficient Risk unchanged controls, conditions, or potential sources of contamination monitoring to warrant 10 pts downgrading risk? **Uses Tablet** Chlorination for treatment Are any 1 Road YES significant (X24) in Risk unchanged contaminant Zone A Reduce risk 1 - 10 pts sources within 8 pts Zone A? Risk posed by potential sources of YES contamination with controls Increase risk 1 - 10 pts 2 pts Existing Risk due to existing 0 pts contamination Are there any NO conditions that Risk unchanged Risk posed by potential sources warrant upgrading Potential of contamination with controls risk? 4 pts Contaminant risks Contaminant Risk YES 4 pts Increase risk 1 - 10 pts 0 pts Contaminant risks* * Truncate risk at 50 pts Contaminant Risk Ratings Risk posed by potential sources of contamination 40 to 50 pts very high = 12 Low 30 to < 40 pts high 20 to < 30 pts medium < 20 pts low







Chart 5. Contaminant risks for Leonard's Landing - Nitrates and Nitrites



Risk Levels for Contaminant Sources identified in Zones A, B and C				
	Zone A	Zones B&C	Total	
Very Highs(s)	0	0	0	
High(s)	0	0	0	
Medium(s)	0	0	0	
Low(s)	1	0	1	

	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	* 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source + 10 pts

Matrix Score	10
--------------	----

Note: Septic systems, sewerlines, and roads are each assigned a risk ranking for each individual contaminant source in the CSI. The VA, however, counts these contaminant sources as a group and assigns a calculated number of either "lows" or "mediums" based on the density.





Chart 5. Contaminant risks for Leonard's Landing - Nitrates and Nitrites Existing NO Are there conditions 3 pts Risk unchanged that warrant upgrading risk? Risk due to existing Potential contamination 12 pts Risk posed by potential sources of contamination with controls Contaminant Risk YES 15 pts Contaminant risks 0 pts Increase risk 1 - 10 pts Risk posed by potential sources of contamination 12 pts Contaminant risks* *Truncate risk at 50 pts 15 Are there sufficient Contaminant Risk Ratings Low controls, conditions, NO Risk unchanged 40 to 50 pts very high or monitoring to warrant downgrading 30 to < 40 ptshigh risk? 20 to < 30 pts medium < 20 pts low YES 0 pts Decrease risk 1 - 10 pts Risk posed by potential sources of contamination with controls 12 pts







Chart 7. Contaminant risks for Leonard's Landing - Volatile Organic Chemicals



Risk Levels for Contaminant Sources identified in Zones A, B and C				
	Zone A	Zones B&C	Total	
Very Highs(s)	0	0	0	
High(s)	0	0	0	
Medium(s)	0	0	0	
Low(s)	1	0	1	

	LOW 10 pts	MEDIUM 20 pts	HIGH 30 pts	VERY HIGH 40 pts
LOW	* 10 sources + 10 pts	≥ 10 sources + 5 pts	≥ 20 sources + 5 pts	
MEDIUM		≥ 2 sources + 5 pts	≥ 5 sources + 5 pts	≥ 10 sources + 5 pts
HIGH			≥ 1 source + 10 pts	≥ 2 sources + 10 pts
VERY HIGH				≥ 1 source + 10 pts

Matrix	Score	10

Note: Septic systems, sewerlines, and roads are each assigned a risk ranking for each individual contaminant source in the CSI. The VA, however, counts these contaminant sources as a group and assigns a calculated number of either "lows" or "mediums" based on the density.

