

I
EIS

Alaska Habitat Management Guide

Life Histories and Habitat Requirements of Fish and Wildlife

Produced by
State of Alaska Department of Fish and Game
Division of Habitat



Juneau, Alaska
1986

*** RECEIVED ***
U. S. COAST GUARD

OCT 15 1986

WATER SAFETY OFFICE
Anchorage, Alaska



Contents

Acknowledgements vii

Introduction 3

Overview of Habitat Management Guides Project 3

Background 3

Purpose 3

Applications 4

Statewide volumes 6

Regional volumes 9

Organization and Use of this Volume 10

Background 10

Organization 10

Species selection criteria 10

Regional Overview Summaries 11

Southwest Region 11

Southcentral Region 13

Arctic Region 15

Western and Interior regions 17

Mammals

Marine Mammals

Belukha whale 21

Bowhead whale 39

Harbor seal 55

Northern fur seal 63

Pacific walrus 71

Polar bear 89

Ringed seal 107

Sea otter 119

Steller sea lion 131

Terrestrial Mammals

Beaver 141

Brown bear 149

Caribou 165

Dall sheep 175

Elk 187

Moose 193

Sitka black-tailed deer 209

Contents (continued)

Birds

Bald Eagle	229
Dabbling ducks	241
Diving ducks	255
Geese	271
Seabirds	291
Trumpeter swan	301
Tundra swan	309

Fish

Freshwater/Anadromous Fish

Arctic char/Dolly Varden	317
Arctic grayling	339
Broad whitefish	361
Burbot	377
Humpback whitefish	389
Lake trout	399
Least cisco	413
Northern pike	425
Rainbow trout/steelhead	439
Salmon	
Chum	457
Chinook	481
Coho	501
Pink	519
Sockeye	537
Sheefish	555

Marine Fish

Arctic cod	571
Capelin	581
Pacific cod	591
Pacific halibut	599
Pacific herring	607
Pacific ocean perch	621
Sablefish	629
Saffron cod	637
Starry flounder	649
Walleye pollock	657
Yelloweye rockfish	667
Yellowfin sole	673

Contents (continued)

Shellfish

- Dungeness crab 683
- King crab 691
- Razor clam 709
- Shrimp 715
- Tanner crab 727

Appendices

- A. Alaska Native Language Names of Selected Fish and Wildlife 743
- B. Directory of Reviewers and Contributors 745
- C. Abbreviations 751
- D. Wildlife Management Goals and Objectives 753
- E. Sea Ice 759

Maps

Introduction

- 1. The six regions of the Alaska Habitat Management Guides 5

Life Histories and Habitat Requirements

Each narrative is preceded by a map of the species' range (map 1 of each narrative).

Pacific walrus

- 1. Range of Pacific walrus 71
- 2. Estimated location of Pacific walrus mating herds, January-March 78
- 3. Estimated maximal extent of calving areas of the Pacific walrus population in mid April and mid June 80

Dabbling ducks

- 1. Range of dabbling ducks 241
- 2. Major coastal marshes of upper Cook Inlet 245

Figures

Introduction

- 1. Types of narratives and maps produced by the AHMG project 7

Tables

Belukha whale

1. Stomach contents of belukha whales collected in Eschschooltz Bay 27
2. Stomach contents of belukha whales collected at Point Hope 28

Bowhead whale

1. Food items from bowhead whale stomach samples collected at Kaktovik, Barter Island, Alaska, Sept.-Oct. 1979 43
2. Quantitative composition of stomach contents from bowhead whales collected at Kaktovik, autumn 1979 45

Brown bear

1. Litter size for brown bears as determined in various studies in North America 157

Sitka black-tailed deer

1. Food species used by Sitka black-tailed deer in the Kodiak and Prince William Sound areas 212

Dabbling ducks

1. Diet composition of 62 dabbling ducks on the west Copper River delta, Sept.-Oct. 1981 249
2. Breeding biology of dabbling ducks 251

Diving ducks

1. Plant and animal species utilized by diving ducks 260
2. Reproductive characteristics of diving ducks 262

Geese

1. Preferred foods and breeding habitat of geese in the Southcentral Region 275
2. Preferred foods and breeding habitat of geese in the Southwest, Western, and Interior regions 276
3. Nesting, rearing, and molting biology of geese in the Southcentral Region 278
4. Nesting, rearing, and molting biology of geese - Southwest, Western, and Interior regions 279
5. Preferred foods and breeding habitat of geese in the Arctic Region 285

Seabirds

1. Seabird life histories 294

Northern pike

1. Food species used by adult northern pike in Alaska 430


Chum salmon

1. Adult chum salmon passage criteria thresholds, for reaches up to 200 ft long within sloughs and side channel habitats in the middle portion of the Susitna River, Alaska 465

Shrimp

1. General characteristics of the five important species of Alaska shrimp 718

Arctic Char Life History and Habitat Requirements
Southwest, Southcentral, Arctic, Western, and Interior Regions



Map 1. Range of arctic char and Dolly Varden in Alaska (ADF&G 1978)

I. NAME

- A. Common Names: Dolly Varden, arctic char
B. Scientific Names: Salvelinus malma (Walbaum), Salvelinus alpinus (Linnaeus)

Dolly Varden and arctic char are two closely related salmonids of the subfamily Salmoninae. Members of the genus Salvelinus are morphologically and ecologically very plastic species. Although this has allowed char to adapt to changing environmental conditions it has been a nightmare for taxonomists, who must attempt to compartmentalize a continuum of life history types and morphological forms into recognizable groups.

Morrow (1980a) stated that both Salvelinus malma and S. alpinus are valid species in Alaska, with S. malma composed of a northern form and a southern form. McPhail (1961) divided the North

American S. alpinus complex into an eastern arctic form and a western arctic-Bering Sea form. The Mackenzie River is the dividing line between the distributions of anadromous populations of the two forms in North America (Craig 1977a), and where the eastern arctic form occurs west of the Mackenzie River it is generally a freshwater lake resident (ibid.). Morrow (1980a) concluded that the northern form of S. malma and the western arctic-Bering Sea form of S. alpinus are identical and should be called S. malma. Within this narrative, we will circumvent the taxonomic problem by referring to all forms of S. malma and S. alpinus jointly as char.

Within each taxonomic form of char there are several different life history patterns. Char may be anadromous, nonanadromous stream residents, or nonanadromous lake residents (Craig 1977a, Morrow 1980a). In the arctic area, there are also a few populations of nonanadromous spring resident fish (Craig 1977a). Lake resident char may be subdivided into dwarf and normal size forms (ibid.). Nonanadromous stream residents may be predominantly dwarf male populations that mate with anadromous females or self-perpetuating populations of male and female fish (Craig 1977a, Morrow 1980a).

C. Native Names: See appendix A.

II. RANGE

A. Statewide

Anadromous and nonanadromous populations are found from the arctic coast south along the western, southwestern, southcentral, and southeastern coastal areas of Alaska. Isolated populations of resident (landlocked) char are found in lakes and streams scattered throughout Interior and Arctic Alaska and on the Kenai Peninsula and Kodiak Island (ADF&G 1978, Morrow 1980b).

B. Regional Distribution Maps

To supplement the distribution information presented in the text, a series of blue-lined reference maps has been prepared for each region. In this series, char distribution information is included on the series of 1:250,000-scale maps titled Distribution of Selected Freshwater Fish. These maps are available for review in ADF&G offices of the region or may be purchased from the contract vendor responsible for their reproduction. In addition, a set of colored 1:1,000,000-scale index maps of selected fish and wildlife species has been prepared and may be found in the Atlas that accompanies each regional guide.

C. Regional Distribution Summary

1. Southwest. Char are widely distributed throughout most systems in the Southwest Region. Important drainages of Bristol Bay include the Togiak River, the Wood River Lakes system, the Tikchik-Nushagak system, the Iliamna-Kvichak system, the Naknek River and Lake, and the Becharof and Ugashik rivers. Some important lake-river systems in the Kodiak region include Uganik, Little River, Karluk, Ayakulik

(Red River), Akalura, Saltery, Buskin, and Barabara lakes. Char are also abundant in the Aleutian Islands (ADF&G 1976, 1977a, and 1977b). (For more detailed narrative information on char distribution in the Southwest Region, see volume 1 of the Alaska Habitat Management Guide for the Southwest Region.)

2. Southcentral. Char are widely distributed throughout the Southcentral area. They are found in the Klutina and Tonsina river drainages and small tributary streams of the Copper River (Williams, pers. comm.). In the Prince William Sound area, nearly all freshwater systems, with the possible exception of short glacial streams on the southeast side of the Kenai Peninsula, contain char. Char are found in lakes and streams on the Kenai Peninsula, most notably the Kenai River, Kasilof River, Deep Creek, Stariski Creek, Anchor River, and lakes in the Swanson River drainages. Char are also found in many streams draining into the west side of Cook Inlet and in the Susitna River drainage (ADF&G 1978). (For more detailed narrative information on char distribution in the Southcentral Region, see volume 2 of the Alaska Habitat Management Guide for the Southcentral Region.)
3. Arctic. Char are found in every major watershed in the Northwest Alaska area, including the Selawik, Kobuk, Noatak, Wulik, and Kivalina rivers (ADF&G 1978, 1984). On the North Slope from Cape Lisburne to Demarcation Point, char are found in most rivers and many lakes. Char have not been reported from rivers or lakes on the arctic coastal plain west of Teshekpuk Lake (Hablett 1979, Bendock and Burr 1984); much of this area, however, has not been extensively surveyed. (For more detailed narrative information on char distribution in the Arctic Region, see volume 2 of the Alaska Habitat Management Guide for the Arctic Region.)
4. Western and Interior. Anadromous char are found south of the Kuskokwim River in Kuskokwim Bay drainages such as the Goodnews, Arolik, and Kanektok rivers (ADF&G 1978, Alt 1977). Dwarf stream resident char and lake resident char are also found in Kuskokwim Bay drainages (Alt 1977). Stream resident char are found in tributaries of the Yukon and Kuskokwim rivers, but they rarely enter the main stem of these rivers (Alt 1980, 1977). Anadromous char are not found in the Yukon or Kuskokwim rivers (ibid.). Lower Kuskokwim tributaries supporting fairly large populations of resident stream char include the Aniak, Kisaralik, Kwithluk, and Tuluksak rivers (ADF&G 1978, Alt 1977). In the Yukon River drainage, char are present as resident species in streams flowing into the Yukon River, from the Andreafsky River upstream to the Melozitna River. The Melozitna River is the furthest upstream where large-size resident char have been found, but dwarf-size resident char are present in small tributary streams further up the Yukon

and in the Upper Tanana drainage (Alt 1981, 1980). (For more detailed narrative information on char distribution in the Western and Interior regions, see Volume 2 of the Alaska Habitat Management Guide for the Western and Interior regions.)

III. PHYSICAL HABITAT REQUIREMENTS

A. Aquatic

Char are found in clear and glacial rivers and lakes, brackish deltas and lagoons (ADF&G 1977a), and nearshore marine waters (Morrow 1980b).

1. Water quality:

a. Temperature. Recorded water temperatures during the spawning period range from 0.5 to nearly 13°C (Morrow 1980b, Moore 1975). In Southeast Alaska, spawning occurs when water temperatures are 5.5 to 6.5°C (Morrow 1980b). Water temperatures on char spawning redds in the Wulik and Kivalina river drainages in Northwest Alaska in August ranged from 6.5 to 11°C (Alt 1978). Egg hatching and alevin development are quite slow but do appear to be dependent upon temperature, with warmer-than-normal temperatures accelerating hatching and resulting in earlier-than-normal fry emergence. Blackett (1968) determined that Southeast Alaska anadromous char eggs held in a hatchery began hatching after 129 days in water with a temperature range of 8.3 to 0.6°C. No upper or lower temperature tolerance limits of char eggs or alevins were found in the literature; however, eggs are frequently exposed to temperatures from 0.0 to 2.2°C during incubation, and Scott and Crossman (1973) report significant egg mortality at temperatures above 7.8°C. Juvenile char have been observed burrowing into the substrate when water temperatures decreased to 2°C (Elliott and Reed 1974). Emigration of char from overwintering areas to summer feeding areas usually occurs after ice breakup in lakes at about 4°C (Armstrong 1965, ADF&G 1977b). Fish reduce feeding and seek overwintering areas when temperatures decrease to or below 5°C (Krueger 1981, ADF&G 1977b). Vertical distribution in lakes appears to be temperature-dependent, with char preferring mid water and bottom depths with temperatures lower than 12.8°C (ADF&G 1976).

b. Dissolved oxygen (D.O). No information was found in the literature on the influence of dissolved oxygen levels on the survival and development of char; however, inferences can be made from work on other salmonid species. Sufficient transport of D.O. to, and metabolic wastes from, developing eggs and alevins by intragravel flow is crucial for survival of eggs and alevins (Vaux

- 1962, Wickett 1958). Relatively low intragravel D.O. levels during the egg development stage may increase egg mortality, influence the rate of egg development, or reduce the fitness of alevins (Alderdice et al. 1958, Silver et al. 1963).
- c. Turbidity. Little work has focused on the influence of turbidity on the survival and development of char; however, inferences can be made from work on other salmonid species. Deposition of fine sediments in spawning areas could reduce the water interchange in the redd and retard or prevent the emergence of fry (Koski 1966). Accumulation of organic debris can reduce dissolved oxygen below safe levels through oxidation (Reiser and Bjornn 1979).
 - d. Salinity. Physiological changes for salinity tolerance of anadromous char probably begin before emigration from freshwater overwintering areas to marine summer feeding areas (Conte and Wagner 1965, Johnson 1980). Roberts (1971), who conducted experiments with a nonanadromous population of char that had been isolated from sea water for about 12,000 years, concluded that nonanadromous char retain a certain degree of salinity tolerance. In the Beaufort Sea in summer, char have been observed in salinities ranging from 2 to 32 parts per thousand (ppt).
2. Water quantity. Sufficient water velocity and depth are required to allow adequate water flow during egg and alevin development. Low flows and cold winter temperatures could cause redds to desiccate or freeze (Krueger 1981). Excessive velocities or flooding can cause egg dislodgement and/or displacement of juvenile (presmolt) char from rearing areas as well as hinder upstream fish migration (ibid.). Juvenile char in the Terror River on Kodiak Island are associated with relatively slow current velocities in pools, quiet side channels, and sloughs and tributaries (Wilson et al. 1981). Habitat preference studies for juvenile char in streams along the route of the Trans Alaska Pipeline System (TAPS) indicated that juvenile char prefer pools with low current velocities, generally below 0.3 m/sec, located adjacent to swift-flowing water (DenBeste and McCart 1984). Anadromous char juveniles and fry in the Sagavanirktok River and neighboring drainages are most abundant in the vicinity of spring water sources (McCart et al. 1972). Proximity to perennial sources of water is probably important in assuring their overwinter survival without having to undertake extensive migrations (ibid.). Char have been observed spawning in water depths of 0.15 to 4.5 m (Krueger 1981, ADF&G 1977b, Alt 1978) and in slow-to-moderate current velocities ranging from 0.3 to 1.2 m/sec (1.0 to 3.8 ft/sec) (Blackett 1968, Scott and Crossman 1973, Alt 1978).

3. Substrate. Preferred spawning substrate is small-to-coarse (walnut-size) gravel (Scott and Crossman 1973, McPhail and Lindsey 1970). Blackett (1968) found char in Southeast Alaska spawning primarily in small gravels, 6 to 50 mm in diameter. Wilson et al. (1981) found char on Kodiak Island spawning on gravels ranging from 2 to 32 mm in diameter. In the Wulik and Kivalina river drainages in Northwest Alaska, char spawn over substrate that is predominantly medium and coarse gravel, with some fine gravel, sand, and boulders (Alt 1978). Char in the Anaktuvuk River also spawn over mixed substrate, ranging from sand to gravel up to three inches (8 cm) in diameter (Bendock 1981). A gravel layer over fertilized eggs in the redd protects eggs from sunlight and predation and reduces disturbance by ice and floods (Krueger 1981). In streams along the route of the TAPS, juvenile char prefer shallow pools with medium-to-coarse rock substrates (DenBeste and McCart 1984). Juvenile char burrow into substrate interstices and logging debris and slash to avoid cooling water temperatures (Elliott and Reed 1974).

B. Cover Requirements

Rocks, logs, root balls, and undercut stream banks in pools, quiet side channels, and high-water overflow areas provide cover for young-of-the-year fish. Char seldom swim near the water surface, preferring to remain near the bottom (Krueger 1981, ADF&G 1977b, ADF&G 1977a). Larger char are most abundant in deeper water in pools, under cutbanks and in quiet water on the downstream side of large boulders (McCart et al. 1972). In-stream vegetation, bank vegetation, shade, in-stream tundra slumps, and rock cover are the most important cover features for juvenile char in streams along the route of the TAPS (DenBeste and McCart 1984). Char overwintering in North Slope streams apparently prefer ice-covered areas even when open-water areas are available (Bendock 1982).

IV. NUTRITIONAL REQUIREMENTS

A. Preferred Foods

Fry begin active feeding as soon as they emerge. Juveniles feed on various winged insects, larvae of mayflies (Ephemeroptera) and midges (Chironomidae), and various small crustaceans (Karzanovskii 1962, Krueger 1981). In the Bristol Bay drainages, fish (sticklebacks, sculpins, blackfish, and salmon fry), fish eggs, and invertebrates (snails, leeches, clams, insects, and insect larvae) are major food sources (Alt 1977, Moriarity 1977, Greenback 1967). Rearing char from the Noatak and Wulik rivers consume fly (Diptera), mayfly, and stonefly (Plecoptera) larvae (DeCicco 1983). Russell (1980) found that char in the Lake Clark area of Bristol Bay consumed gastropods, pelycopods, caddis fly (Trichoptera) larvae and adults, ants and small wasps (Hymenoptera), midge pupae and adults, adult aquatic beetles

(Coleoptera), and small crustaceans (amphipods, copepods, and cladocerans). In the Wood River Lakes system, char feed on sockeye salmon smolt during the smolt's summer migration to Nushagak Bay (Rogers 1972, Buklis 1979). Ruggerone and Rogers (1984) found that the number of smolt consumed in a 24-hour period increased when large smolt migrations occurred. When smolt migrations in excess of 80,000 smolts/24 hours took place, char consumption of smolts increased to 5.6 smolts/char per 24 hours. Palmisano (1971) studied the food habits of char in lakes on Amchitka Island. He found that in lakes with access to the sea, crustaceans and aquatic insects were the major foods. Char in landlocked lakes on Amchitka fed primarily on aquatic insects, fish, and fish eggs (ibid.). Stream resident (nonanadromous) char in the Sagavanirktok and Canning river drainages on the North Slope feed mainly on dipteran larvae (mostly chironomids), stonefly, nymphs, and caddis fly larvae (Craig 1977a, McCart et al. 1972). The diet of large fry in the Canning drainage is generally similar to that of older residents, although the fry feed on fewer kinds and smaller individuals (Craig 1977a). Craig (1977b) found that spring resident char in Sadlerochit Spring in the Arctic National Wildlife Range fed primarily on larvae of stream-dwelling benthic invertebrates, which have high densities in the spring.

In marine waters, smelt, herring, juvenile salmon, sand lance, greenling, sculpins, flounder larvae, and cod are major food components. Amphipods, decapods, mysids, euphausiids, brachiopods, polychaetes, and isopods are also included in their diet (Armstrong and Morrow 1980, Johnson 1980). Townsend (1942) found that char captured near the Shumagin Islands contained large numbers of flounder juveniles and larvae of the sand lance. Off Amchitka Island, char fed on a variety of items, mainly amphipods, mysids, and small fish (Neuhold et al. 1974). Once North Slope char enter coastal waters they feed on insects (chironomid larvae and diptera pupae), crustaceans (Amphipods and mysids), and fish, especially juvenile arctic cod (Boreogadus saida) and fourhorn sculpin (Myoxocephalus quadricornis) (Craig and McCart 1976). Anadromous char feed little while in fresh water. Those that had fed in the Sagavanirktok drainage most frequently consumed arctic char eggs and small arctic char (McCart et al. 1972). Craig and Poulin (1974) found that some char in Weir Creek in the Shaviovik River drainage consumed grayling, and char taken in the Anaktuvuk River during the seaward migration fed on slimy sculpin (Cottus cognatus) and caddis fly larvae (Bendock 1981).

B. Feeding Locations

Juveniles feed primarily from the benthos in low velocity areas along stream and lake margins (Armstrong and Morrow 1980, Johnson 1980). Older char move to deeper and faster stream reaches with higher densities of drifting invertebrates (Krueger 1981, ADF&G 1977b, ADF&G 1977a). Adult anadromous char appear to be equally capable of taking food from mid water or from the bottom (Johnson

1980). Resident char in lakes feed primarily on the lake bottom (Murray, pers. comm.; McCart et al. 1972). McBride (1979) estimated that 40% of the char population in the Wood River Lakes system feed at inlets and outlets of lakes or confluences of rivers and streams during sockeye smolt migrations. Morrow (1980b) states that adult anadromous char consume the majority of their annual diet of small fish and invertebrates in nearshore marine waters.

C. Factors Limiting Availability of Food

Excessive sedimentation may inhibit production of aquatic plants and invertebrate fauna (Hall and McKay 1983) and reduce visual references. While in fresh water, the char may compete directly for food and space with such fishes as grayling, whitefish, sculpins, salmons, and smelt (Armstrong and Morrow 1980). Competitive interactions between char and coho salmon juveniles have been well documented in southeastern Alaska streams (Armstrong 1970, Armstrong and Elliot 1972). Competition for food or space with other species is probably negligible in lakes during the winter (Armstrong and Morrow 1980).

D. Feeding Behavior

Char are carnivorous but have a varied diet, dependent on the size and age of the fish, location, and available food sources. Char may browse along the substrate or consume drifting invertebrates (Armstrong and Elliott 1972). Activity levels and digestive rates drop when freshwater temperatures decrease to or below 5°C (Krueger 1981, ADF&G 1977b). Mature spawners of anadromous populations feed little, if at all, when wintering in fresh water (Morrow 1980b, McCart et al. 1972, Craig 1977a). Stream resident fish feed year-round (McCart et al. 1972, Craig 1977a). When leaving lakes in spring and early summer, char also appear to feed very little (Armstrong 1965).

V. REPRODUCTIVE CHARACTERISTICS

A. Reproductive Habitat

Spawning site selection is influenced by current velocity, water depth, and substrate composition. Spawning sites are usually located in a fairly strong current near the center of the stream in riffles or spring areas at least 0.3 m deep or in gravel-bottomed lakes (Krueger 1981, ADF&G 1977a).

In the Sagavanirktok River and neighboring drainages, all known spawning areas for anadromous char are either in the vicinity of spring sources of mountain streams originating in the Brooks Range or spring-fed tributaries of mountain streams (McCart et al. 1972). Spawning sites in the Anaktuvuk River are also closely associated with spring areas (Bendock 1981). The eggs cannot tolerate freezing, and these are the only stream areas in which winter flow is assured (ibid.). In the Canning River, almost all spawning sites are located in the main channels of the river or in springs originating within or near the Canning floodplain (Craig 1977a). In the Noatak River system, some spawning takes place in

and around springs; however, most spawning occurs downstream of springs in the main channels of streams. The major spawning streams maintain limited flow throughout the winter and are under spring influence for much of their length (DeCicco 1982). Many spawning areas in the Kivalina River are near spring areas; however, in the Wulik River most spawning grounds are not directly influenced by groundwater (Alt 1978).

B. Reproductive Seasonality

All races of char spawn between early July and the beginning of December (Meacham 1977, Alt 1977, DeCicco 1982).

Char have been observed spawning in the Terror and Kizhuyak rivers on Kodiak Island between late August and the end of September (Wilson et al. 1981). On Amchitka Island, Neuhold et al. (1974) observed char spawning from mid October to late November. Char in the Wood River Lakes system spawn in September and October (McBride 1980). Char in the Susitna River drainage also spawn in September and October (ADF&G 1981), and spawning peaks in the Anchor River on the Kenai Peninsula in mid October (Hammarstrom and Wallis 1981) and in Valdez area streams in October and November (Dames and Moore 1979).

The peak of spawning activity in Southeast Alaska occurs between September and November (Blackett 1968, Blackett and Armstrong 1965).

In North Slope rivers, the char spawning period apparently extends over several months; however, the peak of spawning activity probably occurs during late September or early October (McCart et al. 1972, Craig 1977a). In the Anaktuvuk River, spawning has been observed from early September through early November (Bendock 1981). In the Noatak drainage, char are found in the vicinity of the spawning grounds from early July to mid October (DeCicco 1982). In the Noatak and Wulik and Kivalina drainages, char spawners may be divided into two groups: summer spawners that remain in fresh water after overwintering and occupy spawning grounds from early July through mid September, and fall spawners that in-migrate in the fall and spawn from mid September through freeze-up (DeCicco 1982, Alt 1978). In the Wulik and Kivalina rivers, the peaks of spawning appear to be in mid August and mid September (Alt 1978).

Spring resident char in most North Slope streams spawn in November and December (Craig 1977b).

C. Reproductive Behavior

Spawning behavior is similar to that of salmon. Fish are usually paired. The male usually takes no part in the nest-building and spends his time defending the redd from other male spawners. The female excavates the redd, often in typical salmonid fashion by turning on her side and thrashing the substrate with her tail. When the female is ready to deposit her eggs, the pair descend into the redd and press against each other laterally; sperm and eggs are released simultaneously into the redd. After completion of the spawning act, the female may move to the upstream end of

the redd and repeat the digging process, washing gravel downstream over the fertilized eggs. The spawning act may be repeated up to five times; several days are usually required for a female to deposit all her eggs (Morrow 1980b). Morrow (1980b) described the redds as varying from a deep pit to a clean spot on large stones. The dimensions of the redd vary with the size of the female, the substrate, and the current velocities. Male spawners may mate with more than one female; occasionally a female will mate successively with two or more males (Fabricus 1953, Fabricus and Gustafson 1954, Krueger 1981, ADF&G 1977a).

D. Age at Sexual Maturity

Char are an especially slow-growing fish and attain sexual maturity at different ages and sizes, varying with their life history and local environmental conditions. Three life forms of char occur in Alaska: resident lake char, resident stream char, and anadromous char. In general, resident stream char do not grow as large as resident lake or anadromous stream char. Resident stream char commonly occur in dwarf form (sexually mature and fully grown but only six to eight inches in length) (ADF&G 1977a, Russell 1980). Generally, northern populations grow slower, live longer, and reach a smaller maximum size than more southerly populations. Char populations in the south also attain sexual maturity earlier (Morrow 1980b). Males may mature before females. In Kuskokwim Bay drainages, char generally mature at 7 to 10 years (Alt 1977). In the Iliamna system, Metsker (1967) found mature char (life form unknown) as young as four years old. Russell (1980) noted that char in the Lake Clark area apparently become mature at six years of age. Most char in Southeast Alaska reach maturity by ages 4 or 5 (Blackett and Armstrong 1965). Most anadromous char in the Sagavanirktok and Canning river tributaries spawn for the first time at age 7 or 8 (McCart et al. 1972, Craig 1977a). Female char in the Anaktuvuk River mature as early as age 6, males as early as age 7. Stream resident males in these tributaries may mature as early as ages 2 or 3 (ibid.). Most anadromous char from the Wulik and Kivalina rivers in northwest Alaska are mature at age 9 (Alt 1978).

Age at maturity for northern populations of lake resident char is variable, ranging from ages 2 to 9 (Craig 1977a). Males in North Slope spring resident char populations begin maturing at age 2, whereas the youngest mature females are ages 3 or 4 (ibid.). The longevity of char is variable. Char have been found as old as 24 years (Grainger 1953), but most in Southeast Alaska live 8 to 12 years (Armstrong 1963, Heiser 1966, ADF&G 1978).

E. Fecundity

The fecundity of char varies by stock, location, and size of female. Eggs of anadromous stocks are much larger than those of nonanadromous fish and increase in size with fish age and length (Blackett 1968, Morrow 1980b). Fecundity of anadromous char is also higher than for nonanadromous stocks (McCart et al. 1972). In Alaska, the number of eggs generally ranges from 600 to 8,000

per female (ADF&G 1978, Morrow 1980b, McPhail and Lindsey 1970), though Russell (pers. comm.) has observed dwarf, prespawning females with as few as 20 mature eggs in the Tazimina Lakes in Southwest Alaska, and fecundity of female lake resident char from the Canning River drainage ranged from 54 to 1,600 eggs (Craig 1977a).

F. Frequency of Breeding

Though char do suffer a high postspawning mortality rate, a number live to spawn again in subsequent years. Armstrong (1974) found that in a Southeast Alaska population of char 73% spawned once, 26% twice, and 1% three times. Up to 50% of the females spawning for the first time survived to spawn again. In Southeast Alaska, males are much less likely to survive spawning than females. This may also be true of males in North Slope drainages (Armstrong 1974, Yoshihara 1975). Some char spawn in consecutive years; others spawn at two or three-year intervals. Most anadromous char in northern Alaska spawn only every second year (Yoshihara 1975). Craig (1977a), in studies of char in the Canning River, noted that only a small number of char beyond age 9 were found in the population. Because most anadromous char on the North Slope do not mature until ages 7 or 8, relatively few anadromous char may spawn more than twice during their lifetime. Stream resident char, in contrast to the anadromous type, almost always spawn annually (Armstrong and Morrow 1980, Craig 1977a); however, lake resident char from lakes in the Canning River drainage are not annual spawners (Craig 1977a).

G. Incubation Period/Emergence

The time of development varies widely with temperature and stock. Embryo development is slow in cold water temperatures. Eggs incubate over winter, generally four to five months; however, periods of up to eight months have been documented on the North Slope of the Brooks Range (ADF&G 1977a, Yoshihara 1973). The incubation time for char eggs in streambed gravels in the Sagavanirktok and Canning rivers on the North Slope has been estimated to be seven to nine months, though fry in perennial springs may emerge sooner (McCart et al. 1972, Craig 1977a). Eggs hatch as 15-to-20-mm-long alevins (yolk sac fry) in March or April. Yoshihara (1973) observed preemergent fry in the Sagavanirktok River drainage that had probably hatched in April. Alevins remain in the gravel for approximately 18 days while absorbing their yolk sac before they emerge as free-swimming fry (20 to 25 mm) in April to July (ADF&G 1977a, McCart et al. 1972, DeCicco 1982).

In Valdez area streams, fry emerge from the gravel in April and May (Dames and Moore 1979). Peak emergence of fry in the Canning River on the North Slope occurs during the end of May and early June, though fry have been observed in early April (Craig 1977a). In the Noatak drainage, young-of-the-year fry probably emerge sometime in mid July (DeCicco 1982).

VI. MOVEMENTS ASSOCIATED WITH LIFE FUNCTIONS

A. Anadromous

1. Out-migration. Juvenile anadromous char rear in streams and lakes. After a variable number of years (usually at age four or five) most char juveniles undergo a physiological change and, in the spring, migrate seaward as smolts (ADF&G 1977a, 1977b). Most immature and mature char emigrate from overwintering areas to marine summer feeding areas following ice breakup from April to June. In the Sagavanirktok and neighboring North Slope drainages, it is likely that seaward migration occurs during the spring flood in late May and June (McCart et al. 1972). In the Canning River, char begin to leave overwintering areas in May, with a large scale emigration observed in late June and early July (Craig 1977a). In the Anaktuvuk River, char out-migrate in mid June (Bendock 1982). Departure from overwintering habitats may more closely coincide with breakup along the Beaufort Sea coast than with breakup near the overwintering site (ibid.) The char smolt migration in the Anchor River on the Kenai Peninsula takes place in late May and early June (Hammarstrom and Wallis 1983). Nonlake systems may support an additional autumn smolt out-migration (Armstrong 1965 and 1970, Armstrong and Kissner 1969, Dinneford and Elliott 1975, Elliott and Dinneford 1976). In the Noatak, Wulik, and Kivalina rivers, some fish that will spawn in the current year do not migrate seaward in the spring but rather go directly from overwintering to spawning grounds. Some of these summer spawners that spawn in early to mid August probably do migrate to sea for a month or two after spawning (DeCicco 1982, Alt 1978). Alt (1978) estimated that in the Wulik and Kivalina rivers probably in excess of 50% of the prespawning fish do not go out to the ocean at breakup.
2. Marine environment. Individuals remain at sea feeding in the estuary and along the coast for a period of a few weeks to seven months (Morrow 1980b). While in the marine environment, char stay in coastal areas. In some cases char in the Beaufort Sea may travel great distances along the coast (up to 300 km)(Craig and McCart 1976).
3. Spawning and overwintering. Char generally begin reentering fresh water in July and may continue through December, with spawners entering first, followed by immature fish and nonspawners (ADF&G 1977a). Mature anadromous char in the Sagavanirktok River have been taken in the vicinity of spawning tributaries as early as June 26. Mature migrants enter downstream spawning tributaries earlier than those further upstream (McCart et al. 1972). Immature migrants may make this return journey several times before they mature for the first time (ibid.). Although it appears that char return to their natal stream to spawn, nonspawners from Beaufort Sea drainages and from the Wulik and Kivalina river drainages may

overwinter elsewhere in the same drainage or in nonnatal drainages (Craig 1977a, Craig and McCart 1976, DeCicco 1984). Radio-tagging studies of char in the Anaktuvuk River indicate that during the ice-covered months (October to May) char are confined residents of limited overwintering habitat, and movement at this time is minimal (Bendock 1982). In the Chignik River system on the Alaska Peninsula, char, which migrate to sea from April through June, return to Chignik Lake and Black Lake from late July through September to spawn and overwinter (Roos 1959). Emigration of spawned-out char to overwintering areas usually occurs within two weeks after completion of spawning, typically during late October and November. Immature char move to overwintering areas earlier, primarily in July, August, and September (Blackett and Armstrong 1965, Krueger 1981). Adult char usually remain in fresh water through the winter months to avoid the cooler water temperatures of the marine environment (ADF&G 1977a). Overwintering sites include deep lakes, deep river pools, and groundwater spring areas.

B. Nonanadromous

1. Lake residents. Lake resident char move into streams for short periods of time. Studies in the Wood River Lakes system show that discrete subpopulations of resident lake char concentrate at inlets and outlets of the lakes during early summer to feed on out-migrating sockeye salmon smolt (McBride 1979). During late summer, char move to deeper lake waters, probably in response to a declining availability of sockeye salmon smolt and to escape warming surface waters (Nelson 1966). Mature spawners usually move back to the lake margins to spawn in the fall. Resident lake char in the Sagavanirktok drainage do not undertake any long migrations, though they do sometimes enter streams (McCart et al. 1972).
2. Stream residents. Little is known about the life history of resident stream char. In the Sagavanirktok River and neighboring drainages, they are common in mountain and spring streams. Overwintering of stream residents occurs in deep pools of streams and rivers (Morrow 1980b). In Minook Creek, a small tributary of the Yukon River near Rampart, the stream resident char leave smaller tributaries, such as Little Minook Creek and Ruby Creek, in late fall and early winter, apparently spending the winter in the lower reaches of Minook Creek itself (Armstrong and Morrow 1980). During the summer, stream residents in the Kuskokwim and Yukon river drainages congregate in areas where salmon spawn to feed on the salmon eggs (Alt 1980, 1977; ADF&G 1978). In the Sagavanirktok, Canning, Firth, and Babbage rivers, apparently all stream resident char are males;

however, stream resident females have occasionally been found in Fish Creek, Yukon Territory (Craig and McCart 1976). A consequence of this pattern is that female char are significantly more abundant in nearshore areas than are males (ibid.).

VII. FACTORS INFLUENCING POPULATIONS

A. Natural

Natural mortality is largely a result of limited winter habitat. Char that hatch in surface runoff streams must find suitable overwintering areas with open water. Studies in Southeast Alaska indicated that populations of juvenile char suffered 51% mortality in small surface-water streams, versus about 31% mortality in spring-fed streams, from November to June (Elliott and Hubartt 1977). Severe stream flooding can harm developing eggs and embryos and hinder upstream fish migration (Krueger 1981). Low flows and cold winter temperatures could cause redds to desiccate or to freeze. Deposition of fine sediments in the spawning area could retard or prevent fry from emerging (ibid.). Deposition of fine sediments in streams with limited flushing abilities could imbed the substrate material and significantly reduce the available overwintering habitat for juvenile char (Bjornn et al. 1977, Krueger 1981). Postspawning mortality is high and may account for the natural removal of up to 50% of a spawning population (Armstrong and Kissner 1969, ADF&G 1977a). Lake-dwelling populations are often heavily parasitized with nematodes and cestodes (Russell, pers. comm.) There is no significant natural predation on char except for cannibalism (Scott and Crossman 1973, Armstrong and Morrow 1980, Craig 1977a).

B. Human-related

In the Arctic Region, char rely extensively on spring-fed habitats at particular stages in their life cycle. Springs are used as spawning grounds, summer rearing areas of fry and juveniles, and as overwintering areas (Craig 1978). In smaller North Slope drainages, which have few areas suitable for overwintering, it is conceivable that a single spring-fed site might harbor virtually all members of a particular char population, from eggs to mature adults, during the winter period (ibid.). Because stream beds are frozen solid both above and below overwintering sites, char cannot avoid disturbances in their winter habitat (Bendock 1983). Thus any disturbances to spring-fed areas, such as water removal or siltation due to gravel extraction, may have severe deleterious effects on char populations in the entire drainage. The introduction of organic materials under the ice in overwintering areas may reduce dissolved oxygen below the lethal level for any of the life history stages inhabiting the affected stream section (Craig and McCart 1974). A summary of possible impacts from human-related activities includes the following:

- ° Alteration of preferred water temperatures, pH, dissolved oxygen, and chemical composition

- Alteration of preferred water velocity and depth
- Alteration of preferred stream morphology
- Increase in suspended organic or mineral material
- Increase in sedimentation and reduction in permeability of substrate
- Reduction in food supply
- Reduction in protective cover (e.g., overhanging stream banks or vegetation)
- Shock waves in aquatic environment
- Human harvest

(See the Impacts of Land and Water Use volume of this series for additional impacts information.)

VIII. LEGAL STATUS

A. Managerial Authority

The Alaska Board of Fisheries develops regulations governing the sport harvest of fish in Alaska. Research and monitoring of char populations is conducted mainly by the Alaska Department of Fish and Game, Division of Sport Fish.

IX. LIMITATIONS OF INFORMATION

Most life history information on char pertains to the sea-run variety. Little is known about the habits of nonmigratory char. There are very little data relating the various char life stages to the physical and chemical characteristics of their habitats.

REFERENCES

- ADF&G. 1976. A fish and wildlife resource inventory of the Cook Inlet-Kodiak areas. Vol. 2: Fisheries. 434 pp.
- _____. 1978. Alaska's fisheries atlas. Vol. 2 [R.F. Mclean and K.J. Delaney, comps.]. 43 pp. + maps.
- _____. 1981. Susitna Hydroelectric Project Subtask 7.10. Phase 1: final draft report resident fish investigation on the lower Susitna River. Susitna Hydro Aquatic Studies. 138 pp. + appendices.
- _____. 1984. An atlas to the catalog of waters important for spawning, rearing or migration of anadromous fishes. Arctic Region, Resource Management Region V., Div. Habitat, Anchorage. 77 pp. + maps.
- ADF&G, comp. 1977a. A compilation of fish and wildlife resource information for the State of Alaska. Vol. 2: Sport fisheries. 337 pp.
- _____. 1977b. A fish and wildlife resource inventory of the Alaska Peninsula, Aleutian Islands, and Bristol Bay areas. Vol. 2: Fisheries. 556 pp.

Alderdice, D., W. Wickett, and J. Brett. 1958. Some effects of temporary low dissolved oxygen levels on Pacific salmon eggs. J. Fish. Res. Bd. Can. 15(2):229-250.

Alt, K.T. 1977. Inventory and cataloging of sport fish and sport fish waters of Western Alaska. ADF&G, Fed. Aid in Fish Rest. completion rept. Vol. 18. Proj. F-9-9, Job G-I-P.

_____. 1978. Inventory and cataloging of sport fish and sport fish waters of Western Alaska - Wulik-Kivalina rivers study. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 19. Proj. F-9-10, Job G-I-P.

_____. 1980. Inventory and cataloging of sport fish and sport fish waters of Western Alaska. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 21. Proj. F-9-12, Job G-I-P.

_____. 1981. Inventory and cataloging of sport fish and sport fish waters of Western Alaska. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 22. Proj. F-9-13, Job G-I-P.

Armstrong, R. 1963. Investigations of anadromous Dolly Varden population in the Lake Eva-Hanus Bay drainages, Southeastern Alaska. ADF&G, Fed. Aid in Fish Rest. Job completion rept. Vol. 4. Proj. F-5-R-4, Job 3-B. 44 pp.

_____. 1965. Some migratory habits of the anadromous Dolly Varden Salvelinus malma (Walbaum) in Southeast Alaska. ADF&G, Res. Rept. 3:1-26.

_____. 1970. Age, food, and migration of Dolly Varden smolts in Southeast Alaska. J. Fish. Res. Bd. Can. 27:991-1,004.

_____. 1974. Migration of anadromous Dolly Varden (Salvelinus malma) in Southeast Alaska. J. Fish. Res. Bd. Can. 31:435-444. Cited in Armstrong and Morrow 1980.

Armstrong, R., and S. Elliott. 1972. A study of Dolly Varden in Alaska. ADF&G, Fed. Aid in Fish Rest. Ann. prog. rept. Vol. 13. Proj. F-9-4, Job R-IV-D.

Armstrong, R., and P. Kissner. 1969. Investigations of anadromous Dolly Varden populations in Hood Bay, Southeast Alaska. ADF&G, Fed. Aid in Fish Rest. Ann. prog. rept. Vol. 10. Proj. F-9-R-1, Job 2-B.

Armstrong, R.H., and J.E. Morrow. 1980. The Dolly Varden char, Salvelinus malma. Pages 99-140 in E.K. Balon, ed. Chars: salmonid fishes of the genus Salvelinus. The Hauge, the Netherlands: Dr. W. Junk by Publishers. 928 pp.

- Bendock, T.N. 1981. Inventory and cataloging of arctic area waters. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 22. Proj. F-9-13, Job G-I-I.
- _____. 1982. Inventory and cataloging of arctic area waters. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 23. Proj. F-9-14, Job G-I-I.
- _____. 1983. Inventory and cataloging of arctic area waters. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 24. Proj. F-9-15, Job G-I-I.
- Bendock, T.N., and J. Burr. 1984. Inventory and cataloging of arctic area waters. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 25. Proj. F-9-16, Job G-I-I.
- Bjornn, T., M. Brusven, M. Molnau, J. Milligan, R. Klampt, E. Chacho, and C. Schaye. 1977. Transport of granitic sediment in streams and its effects on insects and fish. For Wildlife and Range Experimental Station, completion rept. Water Resource Res. Inst. Proj. B-036-IDA. Univ. Idaho, Moscow. 43 pp.
- Blackett, R.F. 1968. Spawning behavior and early life history of anadromous Dolly Varden, Salvelinus malma (Walbaum), in Southeast Alaska. ADF&G, Res. Rept. 6:1-85
- Blackett R., and R. Armstrong. 1965. Investigations of anadromous Dolly Varden populations in the Lake Eva-Hanus Bay drainages, Southeast Alaska. ADF&G, Div. Sport Fish, Fed. Aid in Fish Rest. Job completion rept. Vol. 6. Proj. F-5-R-6, Job 2-B.
- Buklis, L.S. 1979. Effects of confinement on arctic char in the Wood River Lakes system, Alaska. ADF&G, Informational Leaflet No. 182. 61 pp.
- Conte, F., and H. Wagner. 1965. Development of osmotic and ionic regulation in juvenile steelhead trout, Salmo gairdneri Comp. Biochem. Physiol. 14:603-620.
- Craig, P.C. 1977a. Ecological studies of anadromous and resident populations of arctic char in the Canning River drainage and adjacent coastal waters of the Beaufort Sea, Alaska. Chapter 1 in P. McCart, ed. Fisheries investigations along the North Slope and Beaufort Sea coast in Alaska with emphasis on the arctic char. Canadian Arctic Gas Study Ltd./Alaskan Arctic Gas Study Co. Biol. Rept. Ser., Vol. 41.
- _____. 1977b. Arctic char in Sadlerochit Spring, Arctic National Wildlife Range. Chapter 2 in P. McCart, ed. Fisheries investigations along the North Slope and Beaufort Sea coast in Alaska with emphasis on arctic char. Canadian Arctic Gas Study Ltd./Alaskan Arctic Gas Study Co. Biol. Rept. Ser., Vol. 41.

_____. 1978. Movements of stream-resident and anadromous arctic char (Salvelinus alpinus) in a perennial spring on the Canning River, Alaska. J. Fish. Res. Bd. Can. 35:48-52.

Craig, P.C., and P. McCart. 1974. Fall spawning and overwintering areas of fish populations along routes of proposed pipeline between Prudhoe Bay and the MacKenzie delta 1972-1973. Chapter 3 in P.J. McCart, ed. Fisheries research associated with proposed gas pipeline routes in Alaska, Yukon, and the Northwest Territories. Canadian Arctic Gas Study Co. Biol. Rept. Ser., Vol. 15.

_____. 1976. Fish use of nearshore coastal waters in the western arctic: emphasis on anadromous species. Pages 361-388 in D.W. Hood and D.C. Burrell, eds. Assessment of the arctic marine environment. Occasional Publ. No. 4, Inst. of Mar. Sci., Univ. Alaska, Fairbanks. 468 pp.

Craig, P.C., and V. Poulin. 1974. Life history and movements of grayling (Thymallus arcticus) and juvenile arctic char (Salvelinus alpinus) in a small tundra stream tributary to the Kavik River, Alaska. Chapter 2 in P.J. McCart, ed. Life histories of anadromous and freshwater fish in the western arctic. Canadian Arctic Gas Study Ltd./Alaskan Arctic Gas Study Co. Biol. Rept. Ser., Vol. 20.

Dames and Moore. 1979. Freshwater aquatic habitats of the Valdez area. Pages 263-288 in United States Environmental Protection Agency, environmental impact statement, Alaska petrochemical facility, Valdez, Alaska. Appendix, Vol. 1, EPA 910/9-79-064. 584 pp.

DeCicco, A. 1982. Inventory and cataloging of sport fish and sport fish waters of Western Alaska. Part A: Arctic char life history study. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 23. Proj. F-9-14, Job G-I-P-A.

_____. 1983. Inventory and cataloging of sport fish waters of Western Alaska. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 24. Proj. F-9-15, Job G-I-P-A.

_____. 1984. Inventory and cataloging of sport fish and sport fish waters of western Alaska with emphasis on arctic char life history studies. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 25. Proj. F-9-16, Job G-I-P-A.

DenBeste, J., and P. McCart. 1984. Overview of studies of the long-term effects of the Trans Alaska Pipeline System on fish and aquatic habitats. Vol. 1. Prepared for Alyeska Pipeline Service Co. by Aquatic Environments Inc., Anchorage, AK. 49 pp.

Dinneford, W.B., and S.T. Elliott. 1975. A study of land use activities and their relationship to the sport fish resources in Alaska. ADF&G,

Fed. Aid in Fish Rest. Ann. performance rept. Vol. 16. Proj. F-9-7, Job D-I-B.

- Elliott, S.T., and W.B. Dinneford. 1976. A study of land use activities and their relationship to the sport fish resources in Alaska. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 17. Proj. F-9-8, Job D-I-B. 35 pp.
- Elliott, S.T., and D. Hubartt. 1977. A study of land use activities and their relationship to sport fish resources in Alaska. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 18. Study D-I-B.
- Elliott, S.T., and R.D. Reed. 1974. A study of land-use activities and their relationship to the sportfish resources in Alaska: ecology of rearing fish. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 15. Proj. F-9-6, Job D-I-B.
- Fabricius, E. 1953. Aquarium observations on the spawning behavior of the char, Salmo alpinus. Rept. Inst. Freshwater Res. Drottningholm 35:51-57.d. Cited in Morrow 1980.
- Fabricus, E., and K.J. Gustafson. 1954. Further aquarium observations on the spawning behavior of the char, Salmo alpinus. L. Rept. Inst. Freshwater Res. Drottningholm 35:58-104. Cited in Morrow 1980.
- Grainger, E.H. 1953. On the age, growth, migration, productive potential, and feeding habits of the arctic char (Salvelinus alpinus) of Frosbisher Bay, Baffin Island. J. Fish Res. Bd. Can. 10(6):326-370.
- Greenback, J. 1967. Sport fish survey, Katmai National Monument, Alaska. NMFS, Auke Bay Biological Laboratories Manuscript Rept. 35. 30 pp.
- Hablett, T.R. 1979. Fish inventories conducted within the National Petroleum Reserve on the North Slope of Alaska, 1977-78. Pages 337-406 in National Petroleum Reserve in Alaska Work Group 3. Studies of selected wildlife and fish and their use of habitats on and adjacent to the National Petroleum Reserve in Alaska. Vol. 2. USDI, National Petroleum Reserve in Alaska 105(c) Land Use Study, Anchorage. 423 pp.
- Hall J.E., and D. McKay. 1983. The effects of sedimentation on salmonids and macro-invertebrates - a literature review. ADF&G, Div. Habitat, Anchorage. 31 pp.
- Hammarstrom, S., and J. Wallis. 1981. Inventory and cataloging of Kenai Peninsula and Cook Inlet drainages and fish stocks. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 22. Proj. F-9-13, Job G-I-C.
- _____. 1983. Inventory and cataloging of Kenai Peninsula and Cook Inlet drainages and fish stocks. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 24. Proj. F-9-15, Job G-1-C.

- Heiser, D. 1966. Age and growth of anadromous Dolly Varden char, Salvelinus malma (Walbaum), in Eva Creek, Baranof Island, southeastern Alaska. ADF&G, Research Rept. 5:1-29.
-
- Johnson, L. 1980. The arctic char, Salvelinus alpinus. Pages 15-98 in E.K. Balon, ed. Chars: salmonid fishes of the genus Salvelinus. The Hauge, the Netherlands: Dr. W. Junk by Publishers. 928 pp.
- Karzanovskii, M.Y. 1962. Food of migrating fry of Oncorhynchus gorbuscha and Salvelinus malma in the rivers of Sakhalin. Ryhoe Khoz. 38(6):24-25.
- Koski, K. 1966. The survival of coho salmon (Oncorhynchus kisutch) from egg deposition to emergence in three Oregon coastal streams. M.S. Thesis, Oregon State Univ., Corvallis. 84 pp.
- Krueger, S.W. 1981. Freshwater habitat relationships: Dolly Varden char (Salvelinus malma) (Walbaum). ADF&G, Div. Habitat, Resource Assessment Branch. 38 pp.
- McBride, D.N. 1979. Homing of arctic char, Salvelinus alpinus (Linnaeus), to feeding and spawning sites in the Wood River Lake system, Alaska. M.S. Thesis, Univ. Alaska, Southeastern Senior College, Juneau.
- _____. 1980. Homing of arctic char, Salvelinus alpinus (Linnaeus) to feeding and spawning sites in the Wood River Lake system, Alaska. ADF&G, Informational Leaflet No. 184. 23 pp.
- McCart, P., P. Craig, and H. Bain. 1972. Report on fisheries investigations in the Sagavanirktok River and neighboring drainages. Alyeska Pipeline Service Co. 170 pp.
- McPhail, J.D. 1961. A systematic study of the Salvelinus alpinus complex in North America. J. Fish. Res. Bd. Can. 18:793-816. Cited in Craig 1977.
- McPhail, J.D., and C.C. Lindsey. 1970. Freshwater fishes of northwestern Canada and Alaska. Bull. Fish. Res. Bd. Can. 173:381.
- Meacham, C.P. 1977. Arctic char predations assessment and control investigations within the Wood River System, Alaska, 1975 and 1976. Bristol Bay Data Rept. No. 75. ADF&G, Div. Commer. Fish, Anchorage.
- Metsker, H. 1967. Iliamna Lake watershed freshwater commercial fisheries investigation of 1964. Div. Commer. Fish. Informational Leaflet No. 95. 50 pp.
- Moore, J.W. 1975. Reproductive biology of anadromous arctic char, Salvelinus alpinus (L.), in the Cumberland south area of Baffin Island. J. Fish Biol. 7(2):143-151.

- Moriarty, D.W. 1977. Arctic char in the Wood River Lakes. Final report to Alaska Department of Fish and Game for period November 1, 1975 to October 31, 1976. Univ. Washington, College of Fisheries, Fisheries Research Institute.
- Morrow, J.E. 1980a. Analysis of the Dolly Varden char, Salvelinus malma, of northwestern North America and northeastern Siberia. Pages 323-338 in E.K. Balon, ed. Chars: salmonid fishes of the genus Salvelinus. The Hague, the Netherlands: Dr. W. Junk bv Publishers. 928 pp.
- _____. 1980b. The freshwater fishes of Alaska. Anchorage, AK: Alaska Northwest Publishing Company. 248 pp.
- Murray, J.B. 1984. Personal communications. Area Mgt. Biologist, ADF&G, Div. Sport Fish, Kodiak.
- Nelson, M.O. 1966. Food and distribution of arctic char in Lake Aleknagik, Alaska during the summer of 1962. M.S. Thesis, Univ. Washington. 164 pp. Cited in Moriarty 1977.
- Neuhold, J.M., W.T. Helm, and R.A. Valdez. 1974. Amchitka Bioenvironmental Program: freshwater vertebrate and invertebrate ecology of Amchitka Island. Battelle Memorial Institute, Columbus Laboratories, USAEC Rept. BMI-171-154. 37 pp. Cited in Armstrong and Morrow 1980.
- Palmisano, J.F. 1971. Freshwater food habits of Salvelinus malma (Walbaum) on Amchitka Island, Alaska. M.S. Thesis, Utah State Univ., Logan. 76 pp. Cited in Armstrong and Morrow 1980.
- Reiser, D., and T. Bjornn. 1979. Habitat requirements of anadromous salmonids: influence of forest and rangeland management on anadromous fish habitat in the western United States and Canada. USDA: Forest Service, Gen. Tech. Rept. PNW-96. Pacific Northwest Forest and Range Experimental Station, Portland, OR. 54 pp.
- Roberts, R.A. 1971. Preliminary observations on the ionic regulations of the arctic char Salvelinus alpinus. J. Exp. Biol. 55:213-222. Cited in Johnson 1980.
- Rogers, D.E. 1972. Predator-prey relationship between arctic char and sockeye salmon smolts at the Agulowak River, Lake Aleknagik, in 1971. Fisheries Research Institute, Univ. Washington. Circular No. 72-7.
- Roos J.F. 1959. Feeding habits of the Dolly Varden, Salvelinus malma (Walbaum), at Chignik, Alaska. Trans. Am. Fish. Soc. 88:253-260.
- Ruggerone, G.T., and D.E. Rogers. 1984. Arctic char predation on sockeye salmon smolts at Little Togiak River, Alaska. Fish. Bull. 82:401-410.

Russell, R. 1980. A fisheries inventory of waters in the Lake Clark National Monument Area. ADF&G, Div. Sport Fish, and USDI: NPS. 197 pp.

_____. 1983. Personal communications. Asst. Area Mgt. Biologist, ADF&G, Div. Commer. Fish., King Salmon.

Scott, W.B., and E.J. Crossman. 1973. Freshwater fishes of Canada. Bull. Fish. Res. Bd. Can. 184:965 pp.

Silver, S., C. Warren, and P. Doudoroff. 1963. Dissolved oxygen requirements of developing steelhead trout and chinook salmon embryos at different water velocities. Trans. Am. Fish. Soc. 92(4):327-343.

Townsend, L.D. 1942. The occurrence of flounder post-larvae in fish stomachs. Copeia 1942:126-127. Cited in Armstrong and Morrow 1980.

Vaux, W.A. 1962. Interchange of stream and intragravel water in a salmon spawning riffle. USFWS, Spec. Sci. Rept. Fish No. 405. 11 pp.

Wickett, W. 1958. Review of certain environmental factors affecting the production of pink and chum salmon. J. Fish. Res. Bd. Can. 15(5):1,103-1,126.

Williams, F.T. 1984. Personal communication. Area Mgt. Biologist, ADF&G, Div. Sport Fish, Glennallen.

Wilson, W., E. Trihey, J. Baldrige, C. Evans, J. Thiele, and D. Trudgen. 1981. An assessment of environmental effects of construction and operation of the proposed Terror Lake hydroelectric facility, Kodiak, Alaska. Instream flow studies final report. Arctic Environmental Information and Data Center, Univ. Alaska, Anchorage. 419 pp.

Yoshihara, H.T. 1973. Monitoring and evaluation of arctic waters with emphasis on the North Slope drainages. ADF&G, Fed. Aid in Fish Rest. Ann. prog. rept. Vol. 13. Proj. F-9-5, Job G-III-A.

_____. 1975. Inventory and cataloging of arctic area waters. ADF&G, Fed. Aid in Fish Rest. Ann. performance rept. Vol. 16. Proj. F-9-7, Job G-I-I.